

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide  
Mouse Monoclonal Antibody [Clone NGFR5 + NTR/912 ]  
Catalog # AH12015**

**Specification**

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide  
- Product Information**

Application	,2,3,4,
Primary Accession	<a href="#">P08138</a>
Other Accession	<a href="#">4804</a> , <a href="#">415768</a> , <a href="#">681726</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG's
Calculated MW	75kDa KDa

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide  
- Additional Information**

**Gene ID** 4804

**Other Names**

Tumor necrosis factor receptor superfamily member 16, Gp80-LNGFR, Low affinity neurotrophin receptor p75NTR, Low-affinity nerve growth factor receptor, NGF receptor, p75 ICD, CD271, NGFR, TNFRSF16

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide  
- Protein Information**

**Name** NGFR

**Synonyms** TNFRSF16

**Function**

Low affinity receptor which can bind to NGF, BDNF, NTF3, and NTF4. Forms a heterodimeric receptor with SORCS2 that binds the precursor forms of NGF, BDNF and NTF3 with high affinity, and has much lower affinity for mature NGF and BDNF (PubMed: [24908487](http://www.uniprot.org/citations/24908487)). Plays an important role in differentiation and survival of specific neuronal populations during development (By similarity). Can mediate cell survival as well as cell death of neural cells. Plays a role in the inactivation of RHOA (PubMed: [26646181](http://www.uniprot.org/citations/26646181))

target="\_blank">26646181</a>). Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin- dependent glucose uptake (By similarity). Necessary for the circadian oscillation of the clock genes BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCMgetaN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver (PubMed:<a href="http://www.uniprot.org/citations/23785138" target="\_blank">23785138</a>).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein. Perikaryon {ECO:0000250|UniProtKB:Q9Z0W1}. Cell projection, growth cone {ECO:0000250|UniProtKB:Q9Z0W1}. Cell projection, dendritic spine {ECO:0000250|UniProtKB:Q9Z0W1}

#### NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Melanoma stained with NGFR Monoclonal Antibody (NGFR5 + NTR/912).



Formalin-fixed, paraffin-embedded human Melanoma stained with NGFR Monoclonal Antibody (NGFR5 + NTR/912).

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide - Background**

It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

**NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody - With BSA and Azide - References**

Soland, T.M., et al. 2008. Nerve growth factor receptor (p75NTR) and pattern of invasion predict poor prognosis in oral squamous cell carcinoma. *Histopathology* 53: 62-72. | Wang, W., et al. 2009. Patterns of expression and function of the p75NGFR protein in pancreatic cancer cells and tumours. *Eur. J. Surg. Oncol.* 35: 826-832. |