

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone CCND1/809]
Catalog # AH12238

Specification

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Product Information

Application	,1,2,3,4,
Primary Accession	P24385
Other Accession	595 , 523852 , 667996
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2a, kappa
Calculated MW	36kDa KDa

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 595

Other Names

G1/S-specific cyclin-D1, B-cell lymphoma 1 protein, BCL-1, BCL-1 oncogene, PRAD1 oncogene, CCND1, BCL1, PRAD1

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Protein Information

Name CCND1 {ECO:0000303|PubMed:8204893, ECO:0000312|HGNC:HGNC:1582}

Function

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: 1827756, PubMed: 1833066, PubMed: 19412162, PubMed: 33854235, PubMed: 8114739, PubMed: 8302605). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which

are responsible for the progression through the G(1) phase (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8114739, PubMed:8302605).

Hypophosphorylates RB1 in early G(1) phase (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8114739, PubMed:8302605). Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8302605). Also a substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity (PubMed:15241418). Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (PubMed:9106657). Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner (PubMed:16569215, PubMed:18417529).

Cellular Location


Nucleus. Cytoplasm Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated to the nucleus through interaction with KIP/CIP family members


Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Images







Formalin-fixed, paraffin-embedded human Mantle Cell Lymphoma stained with Cyclin D1 Monoclonal Antibody (CCND1/809).



Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with Cyclin D1 Monoclonal Antibody (CCND1/809).



Flow Cytometry of human Cyclin D1 on HeLa Cells. Black: Cells alone; Green: Isotype Control; Red: PE-labeled Cyclin D1 Monoclonal Antibody (CCND1/809).



Flow Cytometry of human Cyclin D1 on Jurkat Cells. Black: Cells alone; Grey: Isotype Control; Green: AF488-labeled Cyclin D1 Monoclonal Antibody (CCND1/809).

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 36kDa, identified as cyclin D1. Cyclin D1, one of the key cell cycle regulators, is a putative proto-oncogene overexpressed in a wide variety of human neoplasms. This antibody neutralizes the activity of cyclin D1 *in vivo*. About 60% of mantle cell lymphomas (MCL) contain a t(11; 14)(q13; q32) translocation resulting in over-expression of cyclin D1. This antibody is useful in identifying mantle cell lymphomas (cyclin D1 positive) from CLL/SLL and follicular lymphomas (cyclin D1 negative). Occasionally, hairy cell leukemia and plasma cell myeloma weakly express Cyclin D1.

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - References

Baldin V; Lukas J; Marcote MJ; Pagano M; Draetta G. Cyclin D1 is a nuclear protein required for cell cycle progression in G1. *Genes and Development*, 1993, 7(5):812-21