

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide Mouse Monoclonal Antibody [Clone BCL2/782 + BCL2/796] Catalog # AH12254

Specification

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW ,1,2,3,4, <u>P10415</u> <u>596</u>, <u>150749</u> Human Mouse Monoclonal Mouse / IgG's 25-26kDa KDa

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 596

Other Names Apoptosis regulator Bcl-2, BCL2

Storage

Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide - Protein Information

Name BCL2

Function

Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells (PubMed:1508712, PubMed:8183370). Regulates cell death by controlling the mitochondrial membrane permeability (PubMed:11368354). Appears to function in a feedback loop system with caspases (PubMed:11368354). Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1) (PubMed:11368354). Also acts as an inhibitor of autophagy: interacts with BECN1 and AMBRA1 during non-starvation conditions and



inhibits their autophagy function (PubMed:18570871, PubMed:20889974, PubMed:21358617). May attenuate inflammation by impairing NLRP1inflammasome activation, hence CASP1 activation and IL1B release (PubMed:17418785).

Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Nucleus membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P10417}


Tissue Location Expressed in a variety of tissues.

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human non-Hodgkin's Lymphoma stained with Bcl-2 Monoclonal Antibody (BCL2/782 + BCL2/796).





Formalin-fixed, paraffin-embedded human Melanoma stained with Bcl-2 Monoclonal Antibody (BCL2/782 + BCL2/796).

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide -Background

This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 α oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 α oncoprotein inhibits the programmed cell death (apoptosis). In most follicular lymphomas, neoplastic germinal centers express high levels of bcl-2 α protein, whereas the normal or hyperplastic germinal centers are negative. Consequently, this antibody is valuable when distinguishing between reactive and neoplastic follicular proliferation in lymph node biopsies. It may also be used in distinguishing between those follicular lymphomas that express bcl-2 protein and the small number in which the neoplastic cells are bcl-2 negative.

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody - With BSA and Azide -References

Pezzella F et. al.. American Journal of Pathology, 1990, 137(2):225-32. | Adams, J.M., et al. 1998. The Bcl-2 protein family: arbiters of cell survival. Science 281: 1322-1326