

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone 100-1A5]
Catalog # AH12589

Specification

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Product Information

Application	,3,4,
Primary Accession	P29016
Other Accession	910 , 1310
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgM, kappa
Calculated MW	44kDa KDa

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Additional Information

Gene ID 910

Other Names

T-cell surface glycoprotein CD1b, CD1b, CD1B

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Protein Information

Name CD1B

Function

Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Note=Subject to intracellular trafficking between the cell membrane, endosomes and lysosomes.

Tissue Location

Expressed on cortical thymocytes, on certain T-cell leukemias, and in various other tissues

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Images

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - Background

The mouse monoclonal antibody recognizes CD1b, a 44kDa type I glycoprotein associated with beta2-microglobulin (Workshop IV; Code T015). It is expressed on dendritic cells, Langerhans cells, thymocytes, and T acute lymphoblastic leukemia cells. The CD1 multigene family encodes five forms of the CD1 T-cell surface glycoprotein in human, designated CD1A, 1B, 1C, 1D and 1E. CD1, a type 1 membrane protein, has structural similarity to the MHC class I antigen and has been shown to present lipid antigens for recognition by T lymphocytes. Constitutive endocytosis of CD1B molecules and the differential sorting of MHC class II from lysosomes separate peptide- and lipid antigen-presenting molecules during dendritic cell maturation. CD1B is also expressed in interdigitating cells.

CD1b (T-Cell Surface Glycoprotein) Antibody - With BSA and Azide - References

Knapp W. et al. (eds) Leukocyte Typing IV, p251-263, Oxford University Press, Oxford, 1989. | Battistini L, et al. CD1b is expressed in multiple sclerosis lesions. J Neuroimmunol 1996, 67(2):145-151. | Khalili-Shirazi A, et al. The distribution of CD1 molecules in inflammatory neuropathy. J Neurol Sci 1998,158(2):154-163. | Maher JK and Kronenberg M. The role of CD1 molecules in immune responses to infection. Curr Opin Immunol 1997, 9(4):456-461 Blumberg RS et al. Structure and function of the CD1 family of MHC-like cell surface proteins. Immunol Rev 1995, 147:5-29. | Salamone MC et al. Analysis of CD1 molecules on haematological malignancies of myeloid and lymphoid origin. II. Intracellular detection of CD1 antigens. Dis Markers 1990, 8(5):275-281