

**CD3e (T-Cell Marker) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone CRIS-7 ]**  
**Catalog # AH12595**

## Specification

---

### CD3e (T-Cell Marker) Antibody - With BSA and Azide - Product Information

Application	,3,4,
Primary Accession	<a href="#">P07766</a>
Other Accession	<a href="#">916, 3003</a>
Reactivity	Human, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2a, kappa
Calculated MW	20kDa KDa

### CD3e (T-Cell Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 916

#### Other Names

T-cell surface glycoprotein CD3 epsilon chain, T-cell surface antigen T3/Leu-4 epsilon chain, CD3e, CD3E, T3E

#### Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

#### Precautions

CD3e (T-Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

### CD3e (T-Cell Marker) Antibody - With BSA and Azide - Protein Information

Name CD3E

Synonyms T3E

#### Function

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed: <http://www.uniprot.org/citations/2470098> target="\_blank">2470098</a>). In addition of this role of signal transduction in T-cell activation, CD3E plays an essential role in correct T-cell development. Initiates the TCR-CD3 complex assembly by forming the two heterodimers CD3D/CD3E and CD3G/CD3E. Participates also in internalization and cell surface down-regulation of TCR-CD3 complexes via endocytosis sequences

present in CD3E cytosolic region (PubMed:<a href="http://www.uniprot.org/citations/10384095" target="\_blank">10384095</a>, PubMed:<a href="http://www.uniprot.org/citations/26507128" target="\_blank">26507128</a>).

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein

#### **CD3e (T-Cell Marker) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **CD3e (T-Cell Marker) Antibody - With BSA and Azide - Images**

#### **CD3e (T-Cell Marker) Antibody - With BSA and Azide - Background**

Recognizes the  $\epsilon$ -chain of CD3 (Workshop V; Code: CD03.09), which consists of five different polypeptide chains (designated as  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\zeta$ , and  $\eta$ ) with MW ranging from 16-28kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.

#### **CD3e (T-Cell Marker) Antibody - With BSA and Azide - References**

McMichael AJ et al. (eds) Leukocyte Typing III, Oxford University Press, Oxford, 1987. Knapp W et al. (eds) Leukocyte Typing IV, p245 and 1059, Oxford University Press, Oxford, 1989 | Schlossman S et al. (eds) Leukocyte Typing V. Oxford University Press, Oxford, 1995. | Alberola-Ila J et al. Stimulation through the TCR/CD3 complex up-regulates the CD2 surface expression on human T lymphocytes. J Immunol 1991, 146(4):108