

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone 124-1D1]
Catalog # AH12627

Specification

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Product Information

Application	,3,4,
Primary Accession	P09564
Other Accession	924 , 186820
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	40kDa KDa

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 924

Other Names

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protein Information

Name CD7

Function

Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors (PubMed: [7506726](http://www.uniprot.org/citations/7506726)). Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed: [10652336](http://www.uniprot.org/citations/10652336)). In turn, mediates the production of cytokines such as IL-2 (PubMed: [1709867](http://www.uniprot.org/citations/1709867)). On resting NK-cells, CD7 activation results in a significant induction of interferon-gamma levels (PubMed: [7506726](http://www.uniprot.org/citations/7506726)).

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location

Expressed on T-cells and natural killer (NK) cells and their precursors.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Images

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 40kDa, identified as CD7 (Workshop IV; Code T155). CD7 is a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - References

Knapp W et al. eds. Leukocyte typing IV, p341, Oxford University Press, Oxford, 1989 | Miwa H, et al. Biological characteristics of CD7(+) acute leukemia. Leuk. Lymphoma. 1996, 21(3-4):239-244. Rabinowich H, et al. Signaling via CD7 molecules on human NK cells. Induction of tyrosine phosphorylation and beta 1 integrin-mediated adhesion to fibronectin J. Immunol. 1994;153(8):3504-3513. | Saxena A, et al. Biologic and clinical significance of CD7 expression in acute myeloid leukemia. Am J Hematol. 1998, 58(4):278-84