

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone T3-3A1 ]**  
**Catalog # AH12631**

**Specification**

---

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Product Information**

Application	,3,4,
Primary Accession	<a href="#">P09564</a>
Other Accession	<a href="#">924</a> , <a href="#">186820</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	40kDa KDa

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Additional Information**

**Gene ID** 924

**Other Names**

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protein Information**

**Name** CD7

**Function**

Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors (PubMed:<<http://www.uniprot.org/citations/7506726>>7506726</a>). Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed:<<http://www.uniprot.org/citations/10652336>>10652336</a>). In turn, mediates the production of cytokines such as IL-2 (PubMed:<<http://www.uniprot.org/citations/1709867>>1709867</a>). On resting NK-cells, CD7 activation results in a significant induction of interferon-gamma levels (PubMed:<<http://www.uniprot.org/citations/7506726>>7506726</a>).

**Cellular Location**

Membrane; Single-pass type I membrane protein.

**Tissue Location**

Expressed on T-cells and natural killer (NK) cells and their precursors.

## **CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Images**

## **CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - Background**

Recognizes a protein of 40kDa, identified as CD7, a member of the immunoglobulin gene superfamily. Its N-terminal amino acids 1-107 are highly homologous to Ig kappa-L chains whereas the carboxyl-terminal region of the extracellular domain is proline-rich and has been postulated to form a stalk from which the Ig domain projects. CD7 is expressed on the majority of immature and mature T-lymphocytes, and T cell leukemia. It is also found on natural killer cells, a small subpopulation of normal B cells and on malignant B cells. Cross-linking surface CD7 positively modulates T cell and NK cell activity as measured by calcium fluxes, expression of adhesion molecules, cytokine secretion and proliferation. CD7 associates directly with phosphoinositol 3'-kinase. CD7 ligation induces production of D-3 phosphoinositides and tyrosine phosphorylation.

## **CD7 (T-Cell Leukemia Marker) Antibody - With BSA and Azide - References**

Haynes BF, et al. Human lymphocyte antigens: production of a monoclonal antibody that defines functional thymus-derived lymphocyte subsets. Proc. Natl. Acad. Sci. USA 76: 5829-5833, 1979. | Eisenbarth GS, et al. Production of monoclonal antibodies reacting with peripheral blood mononuclear cell surface differentiation antigens. J. Immunol. 124: 1237-1244, 1980. | Haynes BF, et al. Characterization of a monoclonal antibody that defines an immunoregulatory T cell subset for immunoglobulin synthesis in humans. Proc. Natl. Acad. Sci. USA 77: 2914-2918, 1980. | Prieyl JA, LeBien TW. Interleukin 7 independent development of human B cells. Proc. Natl. Acad. Sci. USA 93: 10348-10353, 1996