

**CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone IAP/964 + B6H12.2 ]**  
**Catalog # AH12754**

### Specification

---

**CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Product Information**

Application	,2,3,4,
Primary Accession	<a href="#">Q08722</a>
Other Accession	<a href="#">961</a> , <a href="#">446414</a>
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG's
Calculated MW	~50kDa KDa

**CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Additional Information**

Gene ID 961

**Other Names**

Leukocyte surface antigen CD47, Antigenic surface determinant protein OA3, Integrin-associated protein, IAP, Protein MER6, CD47, CD47, MER6

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Protein Information**

Name CD47

Synonyms MER6

**Function**

Adhesive protein that mediates cell-to-cell interactions (PubMed: [11509594](http://www.uniprot.org/citations/11509594), PubMed: [15383453](http://www.uniprot.org/citations/15383453)). Acts as a receptor for thrombospondin THBS1 and as modulator of integrin signaling through the activation of heterotrimeric G proteins (PubMed: [19004835](http://www.uniprot.org/citations/19004835), PubMed: [7691831](http://www.uniprot.org/citations/7691831), PubMed: [8550562](http://www.uniprot.org/citations/8550562)). Involved in signal transduction, cardiovascular homeostasis,

inflammation, apoptosis, angiogenesis, cellular self-renewal, and immunoregulation (PubMed:<a href="http://www.uniprot.org/citations/11509594" target="\_blank">11509594</a>, PubMed:<a href="http://www.uniprot.org/citations/15383453" target="\_blank">15383453</a>, PubMed:<a href="http://www.uniprot.org/citations/19004835" target="\_blank">19004835</a>, PubMed:<a href="http://www.uniprot.org/citations/27742621" target="\_blank">27742621</a>, PubMed:<a href="http://www.uniprot.org/citations/32679764" target="\_blank">32679764</a>, PubMed:<a href="http://www.uniprot.org/citations/7691831" target="\_blank">7691831</a>, PubMed:<a href="http://www.uniprot.org/citations/8550562" target="\_blank">8550562</a>). Plays a role in modulating pulmonary endothelin EDN1 signaling (PubMed:<a href="http://www.uniprot.org/citations/27742621" target="\_blank">27742621</a>). Modulates nitrous oxide (NO) signaling, in response to THBS1, hence playing a role as a pressor agent, supporting blood pressure (By similarity). Plays an important role in memory formation and synaptic plasticity in the hippocampus (By similarity). Receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells (PubMed:<a href="http://www.uniprot.org/citations/11509594" target="\_blank">11509594</a>). Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation (PubMed:<a href="http://www.uniprot.org/citations/15383453" target="\_blank">15383453</a>). Positively modulates FAS-dependent apoptosis in T-cells, perhaps by enhancing FAS clustering (By similarity). Plays a role in suppressing angiogenesis and may be involved in metabolic dysregulation during normal aging (PubMed:<a href="http://www.uniprot.org/citations/32679764" target="\_blank">32679764</a>). In response to THBS1, negatively modulates wound healing (By similarity). Inhibits stem cell self-renewal, in response to THBS1, probably by regulation of the stem cell transcription factors POU5F1/OCT4, SOX2, MYC/c-Myc and KLF4 (By similarity). May play a role in membrane transport and/or integrin dependent signal transduction (PubMed:<a href="http://www.uniprot.org/citations/7691831" target="\_blank">7691831</a>). May prevent premature elimination of red blood cells (By similarity).

#### Cellular Location

Cell membrane; Multi-pass membrane protein

#### Tissue Location

Very broadly distributed on normal adult tissues, as well as ovarian tumors, being especially abundant in some epithelia and the brain

### CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Images

### CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - Background

This antibody reacts with Ig domain of CD47 protein. CD47, originally named integrin-associated protein (IAP), is a 50kDa protein containing five membrane-spanning sequences and a short cytoplasmic tail. CD47 plays a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It is important in memory formation and

synaptic plasticity in the hippocampus. CD47 may play a role in membrane transport and/or integrin dependent signal transduction.

**CD47 / IAP (Integrin Associated Protein) Antibody - With BSA and Azide - References**

Edris B et. al. Proc Natl Acad Sci USA 109:6656-61 (2012). | Willingham SB et. al. Proc Natl Acad Sci USA 109:6662-7 (2012)