

PITX1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10047

Specification

PITX1 antibody - N-terminal region - Product Information

| | |
|-------------------|--|
| Application | WB |
| Primary Accession | P78337 |
| Other Accession | P78337 , NP_002644 , NM_002653 |
| Reactivity | Human, Mouse, Rat, Pig, Dog, Guinea Pig, Bovine |
| Predicted | Human, Mouse, Rat, Pig, Chicken, Dog, Guinea Pig, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 34 kDa KDa |

PITX1 antibody - N-terminal region - Additional Information

Gene ID 5307

Alias Symbol BFT, POTX, PTX1, CCF

Other Names

Pituitary homeobox 1, Hindlimb-expressed homeobox protein backfoot, Homeobox protein PITX1, Paired-like homeodomain transcription factor 1, PITX1, BFT, PTX1

Target/Specificity

PITX1 is a member of the RIEG/PITX homeobox family, which is involved in organ development and left-right asymmetry. This protein acts as a transcriptional regulator involved in basal and hormone-regulated activity of prolactin. This gene encodes a member of the RIEG/PITX homeobox family, which is in the bicoid class of homeodomain proteins. Members of this family are involved in organ development and left-right asymmetry. This protein acts as a transcriptional regulator involved in basal and hormone-regulated activity of prolactin. Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PITX1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

PITX1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

PITX1 antibody - N-terminal region - Protein Information

Name PITX1

Synonyms BFT, PTX1

Function

Sequence-specific transcription factor that binds gene promoters and activates their transcription. May play a role in the development of anterior structures, and in particular, the brain and facies and in specifying the identity or structure of hindlimb.

Cellular Location


Nucleus.

PITX1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PITX1 antibody - N-terminal region - Images



PITX1 antibody - N-terminal region (AI10047) in Human 293T cells using Western Blot


Host:Rabbit

Target Name:PITX1

Sample Tissue:Human 293T

Antibody Dilution: 1.0µg/ml

PITX1 is supported by BioGPS gene expression data to be expressed in HEK293T




PITX1 antibody - N-terminal region (AI10047) in Human HeLa cells using Western Blot

Host:Rabbit

Target Name:PITX1

Sample Tissue:Human HeLa

Antibody Dilution: 1.0µg/ml PITX1 is strongly supported by BioGPS gene expression data to be expressed in Human HeLa cells



PITX1 antibody - N-terminal region (AI10047) in Human MCF-7 cells using Western Blot

WB Suggested Anti-PITX1 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: MCF7 cell lysate

PITX1 antibody - N-terminal region - Background

This is a rabbit polyclonal antibody against PITX1. It was validated on Western Blot using a cell lysate as a positive control. Abgent strives to provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).