

MAFB antibody - N-terminal region

Rabbit Polyclonal Antibody Catalog # Al10098

Specification

MAFB antibody - N-terminal region - Product Information

Application IHC, WB Primary Accession O9Y5O3

Other Accession <u>09Y503</u>, <u>NP 005452</u>, <u>NM 005461</u>

Reactivity Human, Mouse, Rat, Rabbit, Zebrafish,

Dog, Guinea Pig, Horse, Bovine, Yeast

Predicted Human, Mouse, Rat, Zebrafish, Chicken,

Guinea Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 36 kDa KDa

MAFB antibody - N-terminal region - Additional Information

Gene ID 9935

Alias Symbol KRML, MGC43127

Other Names

Transcription factor MafB, Maf-B, V-maf musculoaponeurotic fibrosarcoma oncogene homolog B, MAFB, KRML

Target/Specificity

MAFB is a basic leucine zipper (bZIP) transcription factor that plays an important role in the regulation of lineage-specific hematopoiesis. The nuclear protein represses ETS1-mediated transcription of erythroid-specific genes in myeloid cells. The protein encoded by this gene is a basic leucine zipper (bZIP) transcription factor that plays an important role in the regulation of lineage-specific hematopoiesis. The encoded nuclear protein represses ETS1-mediated transcription of erythroid-specific genes in myeloid cells. This gene contains no introns.

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-MAFB antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

MAFB antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MAFB antibody - N-terminal region - Protein Information

Name MAFB





Synonyms KRML

Function

Acts as a transcriptional activator or repressor (PubMed:27181683). Plays a pivotal role in regulating lineage-specific hematopoiesis by repressing ETS1-mediated transcription of erythroid- specific genes in myeloid cells. Required for monocytic, macrophage, osteoclast, podocyte and islet beta cell differentiation. Involved in renal tubule survival and F4/80 maturation. Activates the insulin and glucagon promoters. Together with PAX6, transactivates weakly the glucagon gene promoter through the G1 element. SUMO modification controls its transcriptional activity and ability to specify macrophage fate. Binds element G1 on the glucagon promoter (By similarity). Involved either as an oncogene or as a tumor suppressor, depending on the cell context. Required for the transcriptional activation of HOXB3 in the rhombomere r5 in the hindbrain (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00978}.


Tissue Location Ubiquitous..

MAFB antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

MAFB antibody - N-terminal region - Images



MAFB antibody - N-terminal region (Al10098) in Human kidney cells using Immunohistochemistry Human kidney





MAFB antibody - N-terminal region (Al10098) in Human HepG2 cells using Western Blot WB Suggested Anti-MAFB Antibody Titration: 1.25µg/ml

Positive Control: HepG2 cell lysate

MAFB antibody - N-terminal region - Background

This is a rabbit polyclonal antibody against MAFB. It was validated on Western Blot and immunohistochemistry by Abgent. At Abgent we manufacture rabbit polyclonal antibodies on a large scale (200-1000 products/month) of high throughput manner. Our antibodies are peptide based and protein family oriented. We usually provide antibodies covering each member of a whole protein family of your interest. We also use our best efforts to provide you antibodies recognize various epitopes of a target protein. For availability of antibody needed for your experiment, please inquire (sales@abgent.com).