

**DLX3 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10250****Specification****DLX3 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O60479</a>
Other Accession	<a href="#">NM_005220</a> , <a href="#">NP_005211</a>
Reactivity	Human, Mouse, Rat, Pig, Sheep, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Sheep, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

**DLX3 antibody - N-terminal region - Additional Information****Gene ID 1747**

Alias Symbol	TDO, AI4
<b>Other Names</b>	
Homeobox protein DLX-3, DLX3	

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-DLX3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

DLX3 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**DLX3 antibody - N-terminal region - Protein Information****Name DLX3****Function**

Transcriptional activator (By similarity). Activates transcription of GNRHR, via binding to the downstream activin regulatory element (DARE) in the gene promoter (By similarity).

**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}. Cytoplasm {ECO:0000250|UniProtKB:Q64205}

## **DLX3 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)