

ZBTB20 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI10280**Specification****ZBTB20 antibody - middle region - Product Information**

Application	WB
Primary Accession	O9HC78
Other Accession	NM_015642 , NP_056457
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	73kDa KDa

ZBTB20 antibody - middle region - Additional Information**Gene ID** 26137**Alias Symbol** HOF, DPZF, ODA-8S, ZNF288**Other Names**

Zinc finger and BTB domain-containing protein 20, Dendritic-derived BTB/POZ zinc finger protein, Zinc finger protein 288, ZBTB20, DPZF, ZNF288

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZBTB20 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZBTB20 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB20 antibody - middle region - Protein Information**Name** ZBTB20**Synonyms** DPZF, ZNF288**Function**May be a transcription factor that may be involved in hematopoiesis, oncogenesis, and immune responses (PubMed: <http://www.uniprot.org/citations/11352661> target="_blank">11352661). Plays a role in postnatal myogenesis, may be involved in the regulation of satellite cells self-renewal (By similarity).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q8K0L9}.

Tissue Location

Expressed in spleen, lymph node, thymus, peripheral blood leukocytes, and fetal liver.

ZBTB20 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

