

POGK Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI10445

Specification

POGK Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	G3V1P0
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

POGK Antibody - C-terminal region - Additional Information

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-POGK antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

POGK Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

POGK Antibody - C-terminal region - Protein Information

POGK Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

POGK Antibody - C-terminal region - Images



90 kDa__
65 kDa__
40 kDa__
29 kDa__
22 kDa__

Host:Rabbit

Target Name:POGK

Sample Tissue: Large Intestine Tumor lysates

Antibody Dilution:1.µg/ml