

**STAT1 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI10505**

**Specification**

**STAT1 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">P42224</a>
Other Accession	<a href="#">NM_007315</a> , <a href="#">NP_009330</a>
Reactivity	Human, Mouse, Rat, Pig, Sheep, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Pig, Sheep, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87kDa KDa

**STAT1 antibody - N-terminal region - Additional Information**

**Gene ID 6772**

Alias Symbol **DKFZp686B04100, ISGF-3, STAT91, CANDF7**

**Other Names**

Signal transducer and activator of transcription 1-alpha/beta, Transcription factor ISGF-3 components p91/p84, STAT1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-STAT1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

STAT1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**STAT1 antibody - N-terminal region - Protein Information**

**Name STAT1**

**Function**

Signal transducer and transcription activator that mediates cellular responses to interferons (IFNs), cytokine KITLG/SCF and other cytokines and other growth factors (PubMed:<a href="http://www.uniprot.org/citations/12764129" target="\_blank">12764129</a>, PubMed:<a href="http://www.uniprot.org/citations/12855578" target="\_blank">12855578</a>, PubMed:<a href="http://www.uniprot.org/citations/15322115" target="\_blank">15322115</a>, PubMed:<a

href="http://www.uniprot.org/citations/23940278" target="\_blank">23940278</a>, PubMed:<a href="http://www.uniprot.org/citations/34508746" target="\_blank">34508746</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>, PubMed:<a href="http://www.uniprot.org/citations/9724754" target="\_blank">9724754</a>). Following type I IFN (IFN-alpha and IFN-beta) binding to cell surface receptors, signaling via protein kinases leads to activation of Jak kinases (TYK2 and JAK1) and to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize and associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus (PubMed:<a href="http://www.uniprot.org/citations/28753426" target="\_blank">28753426</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>). ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IFN-stimulated genes (ISG), which drive the cell in an antiviral state (PubMed:<a href="http://www.uniprot.org/citations/28753426" target="\_blank">28753426</a>, PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>). In response to type II IFN (IFN-gamma), STAT1 is tyrosine- and serine-phosphorylated (PubMed:<a href="http://www.uniprot.org/citations/26479788" target="\_blank">26479788</a>). It then forms a homodimer termed IFN-gamma-activated factor (GAF), migrates into the nucleus and binds to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state (PubMed:<a href="http://www.uniprot.org/citations/8156998" target="\_blank">8156998</a>). Becomes activated in response to KITLG/SCF and KIT signaling (PubMed:<a href="http://www.uniprot.org/citations/15526160" target="\_blank">15526160</a>). May mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed:<a href="http://www.uniprot.org/citations/19088846" target="\_blank">19088846</a>). Involved in food tolerance in small intestine: associates with the Gasdermin-D, p13 cleavage product (13 kDa GSDMD) and promotes transcription of CIITA, inducing type 1 regulatory T (Tr1) cells in upper small intestine (By similarity).

#### Cellular Location

Cytoplasm. Nucleus Note=Translocated into the nucleus upon tyrosine phosphorylation and dimerization, in response to IFN-gamma and signaling by activated FGFR1, FGFR2, FGFR3 or FGFR4 (PubMed:15322115). Monomethylation at Lys- 525 is required for phosphorylation at Tyr-701 and translocation into the nucleus (PubMed:28753426). Translocates into the nucleus in response to interferon-beta stimulation (PubMed:26479788)

#### STAT1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

