

NR5A1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11228**Specification**

NR5A1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P33242
Other Accession	NM_139051 , NP_620639
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Dog
Predicted Host	Human, Rabbit
Clonality	Rabbit
Calculated MW	Polyclonal 51kDa KDa

NR5A1 antibody - C-terminal region - Additional Information**Gene ID** 26423**Alias Symbol** ELP, SF1, SF-1, Ad4BP, ELP-3, Ftzf1, Ftz-F1**Other Names**

Steroidogenic factor 1, SF-1, STF-1, Adrenal 4-binding protein, Embryonal LTR-binding protein, ELP, Embryonal long terminal repeat-binding protein, Fushi tarazu factor homolog 1, Nuclear receptor subfamily 5 group A member 1, Steroid hormone receptor Ad4BP, Steroid hydroxylase positive regulator, Nr5a1, Ftzf1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NR5A1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

NR5A1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NR5A1 antibody - C-terminal region - Protein Information**Name** Nr5a1**Synonyms** Ftzf1**Function**

Transcriptional activator. Seems to be essential for sexual differentiation and formation of the primary steroidogenic tissues. Binds to the Ad4 site found in the promoter region of steroidogenic P450 genes such as CYP11A, CYP11B and CYP21B. Also regulates the AMH/Muellerian inhibiting

substance gene as well as the AHCH and STAR genes. 5'-YCAAGGYC-3' and 5'-RRAGGTCA-3' are the consensus sequences for the recognition by NR5A1. The SFPQ-NONO-NR5A1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity (By similarity). Transcription repressor of the Moloney leukemia virus long terminal repeat in undifferentiated murine embryonal carcinoma cells. Binds phosphatidylcholine and phospholipids with a phosphatidylinositol (PI) headgroup, in particular phosphatidyl(3,4)bisphosphate, phosphatidyl(3,5)bisphosphate and phosphatidyl(3,4,5)triphosphate. Activated by the phosphorylation of NR5A1 by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation.

Cellular Location


Nucleus.

NR5A1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR5A1 antibody - C-terminal region - Images



WB Suggested Anti-NR5A1 Antibody Titration: 0.3125 μ g/ml

ELISA Titer: 1:312500

Positive Control: Human Adrenal gland

NR5A1 antibody - C-terminal region - References

Fatchiyah, et al., (2006) Biochem. Biophys. Res. Commun. 341 (4), 1036-1045
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.