

TRIM21 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11383**Specification**

TRIM21 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P19474
Other Accession	NM_003141 , NP_003132
Reactivity	Human, Rat, Zebrafish, Pig, Horse, Bovine
Predicted	Human, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

TRIM21 antibody - C-terminal region - Additional Information**Gene ID 6737**Alias Symbol **RNF81, RO52, SSA, SSA1****Other Names**

E3 ubiquitin-protein ligase TRIM21, 6.3.2.-, 52 kDa Ro protein, 52 kDa ribonucleoprotein autoantigen Ro/SS-A, RING finger protein 81, Ro(SS-A), Sjogren syndrome type A antigen, SS-A, Tripartite motif-containing protein 21, TRIM21, RNF81, RO52, SSA1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-TRIM21 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TRIM21 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TRIM21 antibody - C-terminal region - Protein InformationName TRIM21 ([HGNC:11312](#))

Synonyms RNF81, RO52, SSA1

FunctionE3 ubiquitin-protein ligase whose activity is dependent on E2 enzymes, UBE2D1, UBE2D2, UBE2E1 and UBE2E2 (PubMed: <http://www.uniprot.org/citations/16297862> target="_blank">16297862, PubMed: <http://www.uniprot.org/citations/16316627> target="_blank">16316627, PubMed: <http://www.uniprot.org/citations/16472766> target="_blank">16472766, PubMed: <http://www.uniprot.org/citations/16880511>

target="_blank">16880511, PubMed:18022694, PubMed:18361920, PubMed:18641315, PubMed:18845142, PubMed:19675099, PubMed:26347139). Forms a ubiquitin ligase complex in cooperation with the E2 UBE2D2 that is used not only for the ubiquitination of USP4 and IKBKB but also for its self-ubiquitination (PubMed:16880511, PubMed:19675099). Component of cullin-RING-based SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complexes such as SCF(SKP2)-like complexes (PubMed:16880511). A TRIM21-containing SCF(SKP2)-like complex is shown to mediate ubiquitination of CDKN1B ('Thr-187' phosphorylated- form), thereby promoting its degradation by the proteasome (PubMed:16880511). Monoubiquitinates IKBKB that will negatively regulates Tax-induced NF-kappa-B signaling (PubMed:19675099). Negatively regulates IFN-beta production post-pathogen recognition by catalyzing polyubiquitin-mediated degradation of IRF3 (PubMed:18641315). Mediates the ubiquitin-mediated proteasomal degradation of IgG1 heavy chain, which is linked to the VCP-mediated ER-associated degradation (ERAD) pathway (PubMed:18022694). Promotes IRF8 ubiquitination, which enhanced the ability of IRF8 to stimulate cytokine genes transcription in macrophages (By similarity). Plays a role in the regulation of the cell cycle progression (PubMed:16880511). Enhances the decapping activity of DCP2 (PubMed:18361920). Exists as a ribonucleoprotein particle present in all mammalian cells studied and composed of a single polypeptide and one of four small RNA molecules (PubMed:1985094, PubMed:8666824). At least two isoforms are present in nucleated and red blood cells, and tissue specific differences in RO/SSA proteins have been identified (PubMed:8666824). The common feature of these proteins is their ability to bind HY RNAs.2 (PubMed:8666824). Involved in the regulation of innate immunity and the inflammatory response in response to IFNG/IFN-gamma (PubMed:26347139). Organizes autophagic machinery by serving as a platform for the assembly of ULK1, Beclin 1/BECN1 and ATG8 family members and recognizes specific autophagy targets, thus coordinating target recognition with assembly of the autophagic apparatus and initiation of autophagy (PubMed:26347139). Regulates also autophagy through FIP200/RB1CC1 ubiquitination and subsequent decreased protein stability (PubMed:36359729). Represses the innate antiviral response by facilitating the formation of the NMI-IFI35 complex through 'Lys-63'- linked ubiquitination of NMI (PubMed:26342464). During viral infection, promotes cell pyroptosis by mediating 'Lys-6'-linked ubiquitination of ISG12a/IFI27, facilitating its translocation into the mitochondria and subsequent CASP3 activation (PubMed:36426955). When up-regulated through the IFN/JAK/STAT signaling pathway, promotes 'Lys-27'-linked ubiquitination of MAVS, leading to the recruitment of TBK1 and up- regulation of innate immunity (PubMed:29743353). Mediates 'Lys-63'- linked polyubiquitination of G3BP1 in response to heat shock, leading to stress granule disassembly (PubMed:36692217).

target="_blank">36692217).

Cellular Location

Cytoplasm. Cytoplasmic vesicle, autophagosome. Nucleus. Cytoplasm, P-body. Cytoplasm, Stress granule. Note=Enters the nucleus upon exposure to nitric oxide (PubMed:18361920). Localizes to small dot- or rod-like structures in the cytoplasm, called processing bodies (P-bodies) that are located underneath the plasma membrane and also diffusely in the cytoplasm (PubMed:18361920). They are located along the microtubules and are highly motile in cells (PubMed:18361920). Colocalizes with DCP2 in P-bodies (PubMed:18361920). Localizes to stress granules in response to oxidative stress (PubMed:36692217).

Tissue Location


Isoform 1 and isoform 2 are expressed in fetal and adult heart and fetal lung

TRIM21 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIM21 antibody - C-terminal region - Images



WB Suggested Anti-TRIM21 Antibody Titration: 1.25µg/ml
Positive Control: Transfected 293T

TRIM21 antibody - C-terminal region - References

Yamochi,T., (2008) Biochem. Biophys. Res. Commun. 370 (1), 195-199 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.