

ZNF326 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11668**Specification**

ZNF326 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q5BKZ1
Other Accession	NM_182976 , NP_892021
Reactivity	Human, Rat, Dog
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66kDa KDa

ZNF326 antibody - C-terminal region - Additional Information**Gene ID** 284695**Alias Symbol** **FLJ20403, MGC61591, ZAN75, Zfp326, dj871E2.1****Other Names**

DBIRD complex subunit ZNF326, Zinc finger protein 326, Zinc finger protein interacting with mRNPs and DBC1, ZNF326, ZIRD

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ZNF326 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF326 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF326 antibody - C-terminal region - Protein Information**Name** ZNF326**Synonyms** ZIRD**Function**

Core component of the DBIRD complex, a multiprotein complex that acts at the interface between core mRNP particles and RNA polymerase II (RNAPII) and integrates transcript elongation with the regulation of alternative splicing; the DBIRD complex affects local transcript elongation rates and alternative splicing of a large set of exons embedded in (A + T)-rich DNA regions. May play a role in neuronal differentiation and is able to bind DNA and activate expression in vitro.


Cellular Location

Nucleus matrix.


ZNF326 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF326 antibody - C-terminal region - Images

Human kidney



Human Liver



70 kDa
60 kDa
48 kDa
36 kDa
21 kDa

WB Suggested Anti-ZNF326 Antibody Titration: 0.625 μ g/ml
Positive Control: Jurkat cell lysate

ZNF326 antibody - C-terminal region - References

Strausberg, R.L., (2002) Proc. Natl. Acad. Sci. U.S.A. 99 (26), 16899-16903
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.