

HNRPK antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11741**Specification**

HNRPK antibody - N-terminal region - Product Information

| | |
|-------------------|--|
| Application | IHC, WB |
| Primary Accession | P61978 |
| Other Accession | NM_031262 , NP_112552 |
| Reactivity | Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Horse, Bovine, Dog |
| Predicted | Human, Mouse, Rat, Rabbit, Zebrafish, Chicken, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 51kDa KDa |

HNRPK antibody - N-terminal region - Additional Information**Gene ID** 3190**Alias Symbol** CSBP, TUNP, HNRPK**Other Names**

Heterogeneous nuclear ribonucleoprotein K, hnRNP K, Transformation up-regulated nuclear protein, TUNP, HNRNPK, HNRPK

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-HNRPK antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

HNRPK antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HNRPK antibody - N-terminal region - Protein Information**Name** HNRNPK**Synonyms** HNRPK**Function**

One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA. Plays an important role in p53/TP53 response to DNA damage, acting at the level of both transcription activation and

repression. When sumoylated, acts as a transcriptional coactivator of p53/TP53, playing a role in p21/CDKN1A and 14-3-3 sigma/SFN induction (By similarity). As far as transcription repression is concerned, acts by interacting with long intergenic RNA p21 (lincRNA-p21), a non-coding RNA induced by p53/TP53. This interaction is necessary for the induction of apoptosis, but not cell cycle arrest. As part of a ribonucleoprotein complex composed at least of ZNF827, HNRNPL and the circular RNA circZNF827 that nucleates the complex on chromatin, may negatively regulate the transcription of genes involved in neuronal differentiation (PubMed:33174841).

Cellular Location


Cytoplasm. Nucleus, nucleoplasm. Cell projection, podosome. Note=Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear (PubMed:18775702)

HNRPK antibody - N-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HNRPK antibody - N-terminal region - Images



Rabbit Anti-HNRPK Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-HNRPK Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: HepG2 cell lysate

HNRPK antibody - N-terminal region - References

Nakagawa, J., (2006) Br. J. Cancer 94 (4), 586-592
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.