

**CHST1 antibody - middle region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12584****Specification**

---

**CHST1 antibody - middle region - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O43916</a>
Other Accession	<a href="#">NM_003654</a> , <a href="#">NP_003645</a>
Reactivity	<b>Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Bovine, Guinea Pig, Dog</b>
Predicted	<b>Human, Mouse, Rat, Rabbit, Zebrafish, Chicken, Bovine, Dog</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>47kDa KDa</b>

**CHST1 antibody - middle region - Additional Information****Gene ID** 8534**Alias Symbol** **C6ST, KS6ST, KSGal6ST, KSST, GST-1****Other Names**

Carbohydrate sulfotransferase 1, 2.8.2.21, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 1, GST-1, Keratan sulfate Gal-6 sulfotransferase, KS6ST, KSGal6ST, KSST, CHST1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-CHST1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CHST1 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

**CHST1 antibody - middle region - Protein Information****Name** CHST1 ([HGNC:1969](#))**Function**

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of internal galactose (Gal) residues of keratan. Cooperates with B4GALT4 and B3GNT7 glycosyltransferases and CHST6 sulfotransferase to construct and elongate disulfated disaccharide unit [-&gt;3(6-sulfoGalbeta)1-&gt;4(6-sulfoGlcNAcbeta)1-&gt;] within keratan sulfate polymer (PubMed:&lt;a href="http://www.uniprot.org/citations/10642612"&gt;

target="\_blank">10642612</a>, PubMed:<a href="http://www.uniprot.org/citations/17690104" target="\_blank">17690104</a>, PubMed:<a href="http://www.uniprot.org/citations/9405439" target="\_blank">9405439</a>). Has a preference for sulfating keratan sulfate, but it also transfers sulfate to the unsulfated polymer (PubMed:<a href="http://www.uniprot.org/citations/9405439" target="\_blank">9405439</a>). Involved in biosynthesis of phosphacan, a major keratan sulfate proteoglycan in the developing brain (By similarity). Involved in biosynthesis of 6-sulfoGalbeta- containing O-linked glycans in high endothelial venules of lymph nodes. May act in a synergistic manner with CHST4 to generate sialyl 6',6- disulfo Lewis X motif, a recognition determinant for immune cell receptors implicated in leukocyte trafficking (PubMed:<a href="http://www.uniprot.org/citations/10330415" target="\_blank">10330415</a>). Catalyzes sulfation of N-acetyllactosamine (LacNAc) oligosaccharides with highest efficiency for sialylated LacNAc structures (PubMed:<a href="http://www.uniprot.org/citations/10642612" target="\_blank">10642612</a>).

### Cellular Location

Golgi apparatus membrane; Single- pass type II membrane protein

### Tissue Location


Widely expressed at low level. Expressed in brain and skeletal muscle. Expressed by high endothelial cells (HEVs) and leukocytes.

## CHST1 antibody - middle region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## CHST1 antibody - middle region - Images



WB Suggested Anti-CHST1 Antibody Titration: 0.2-1 µg/ml  
Positive Control: Human kidney



Host:Rabbit  
Target Name:CHST1  
Sample Tissue:Human Fetal Heart  
Antibody Dilution: 1.0µg/ml



Host:Rabbit  
Target Name:CHST1  
Sample Tissue:Human Fetal Lung  
Antibody Dilution: 1.0µg/ml

### CHST1 antibody - middle region - References

Kitayama,K.,(2007).*J.Biol.Chem.*282(41),30085-30096  
ReconstitutionandStorage:Forshorttermuse,storeat2-8°Cupto1week.Forlongtermstorage,storeat-20°Cinsmallaliquotstopreventfreeze-thawcycles.