

**FMNL2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12831****Specification**

---

**FMNL2 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O96PY5</a>
Other Accession	<a href="#">NM_052905</a> , <a href="#">NP_443137</a>
Reactivity	Human, Mouse, Rat, Guinea Pig
Predicted	Human, Mouse, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	124kDa KDa

**FMNL2 antibody - N-terminal region - Additional Information****Gene ID** 114793**Alias Symbol** FHOD2, FLJ37546**Other Names**

Formin-like protein 2, Formin homology 2 domain-containing protein 2, FMNL2, FHOD2, KIAA1902

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-FMNL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

FMNL2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**FMNL2 antibody - N-terminal region - Protein Information****Name** FMNL2 ([HGNC:18267](#))**Synonyms** FHOD2, KIAA1902**Function**

Plays a role in the regulation of cell morphology and cytoskeletal organization. Required in the cortical actin filament dynamics.

**Cellular Location**


Cytoplasm.

## FMNL2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## FMNL2 antibody - N-terminal region - Images



WB Suggested Anti-FMNL2 Antibody Titration: 0.2-1  $\mu$ g/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

## FMNL2 antibody - N-terminal region - References

Colland, F., (2004) *Genome Res.* 14(7), 1324-1332. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.