

CAND2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13189**Specification**

CAND2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O75155
Other Accession	NM_012298 , NP_036430
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	123kDa KDa

CAND2 antibody - N-terminal region - Additional Information**Gene ID** 23066**Alias Symbol** KIAA0667, TIP120B, Tp120b**Other Names**

Cullin-associated NEDD8-dissociated protein 2, Cullin-associated and neddylation-dissociated protein 2, Epididymis tissue protein Li 169, TBP-interacting protein of 120 kDa B, TBP-interacting protein 120B, p120 CAND2, CAND2, KIAA0667, TIP120B

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CAND2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CAND2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CAND2 antibody - N-terminal region - Protein Information**Name** CAND2**Synonyms** KIAA0667, TIP120B**Function**

Probable assembly factor of SCF (SKP1-CUL1-F-box protein) E3 ubiquitin ligase complexes that promotes the exchange of the substrate- recognition F-box subunit in SCF complexes, thereby playing a key role in the cellular repertoire of SCF complexes.

Cellular Location

Nucleus.


Tissue Location

Expressed in epididymis (at protein level).

CAND2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CAND2 antibody - N-terminal region - Images

WB Suggested Anti-CAND2 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: Human Muscle

CAND2 antibody - N-terminal region - References

- Li J., et al. Mol. Cell. Proteomics 9:2517-2528(2010).
Muzny D.M., et al. Nature 440:1194-1198(2006).
Bienvenut W.V., et al. Submitted (MAR-2009) to UniProtKB.
Bienvenut W.V., et al. Submitted (MAR-2009) to UniProtKB.
Ishikawa K., et al. DNA Res. 5:169-176(1998).