

CRBN antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14025**Specification**

CRBN antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O96SW2
Other Accession	NM_016302 , NP_057386
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50kDa KDa

CRBN antibody - N-terminal region - Additional Information**Gene ID** 51185**Alias Symbol** [DKFZp781K0715](#), [MGC27358](#), [MRT2A](#), [MRT2](#)
Other Names
Protein cereblon, CRBN**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CRBN antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CRBN antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CRBN antibody - N-terminal region - Protein Information**Name** CRBN**Function**Substrate recognition component of a DCX (DDB1-CUL4-X-box) E3 protein ligase complex that mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as MEIS2, ILF2 or GLUL (PubMed: [26990986](http://www.uniprot.org/citations/26990986), PubMed: [33009960](http://www.uniprot.org/citations/33009960)). Normal degradation of key regulatory proteins is required for normal limb outgrowth and expression of the fibroblast growth factor FGF8 (PubMed: [20223979](http://www.uniprot.org/citations/20223979), PubMed: [20223979](#)).

<http://www.uniprot.org/citations/24328678> target="_blank">24328678, PubMed:25043012, PubMed:25108355). Maintains presynaptic glutamate release and consequently cognitive functions, such as memory and learning, by negatively regulating large-conductance calcium-activated potassium (BK) channels in excitatory neurons (PubMed:18414909, PubMed:29530986). Likely to function by regulating the assembly and neuronal surface expression of BK channels via its interaction with KCNT1 (PubMed:18414909). May also be involved in regulating anxiety-like behaviors via a BK channel-independent mechanism (By similarity). Plays a negative role in TLR4 signaling by interacting with TRAF6 and ECSIT, leading to inhibition of ECSIT ubiquitination, an important step of the signaling (PubMed:31620128).

Cellular Location

Cytoplasm. Nucleus. Membrane; Peripheral membrane protein

Tissue Location


Widely expressed. Highly expressed in brain.

CRBN antibody - N-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRBN antibody - N-terminal region - Images



WB Suggested Anti-CRBN Antibody Titration: 0.2-1 µg/ml
Positive Control: COLO205 cell lysate



Host: Rabbit
Target Name: CRBN
Sample Tissue: RPMI-8226 Whole Cell lysates
Antibody Dilution: 0.5µg/ml

CRBN antibody - N-terminal region - References

- Ota T., et al. Nat. Genet. 36:40-45(2004).
Muzny D.M., et al. Nature 440:1194-1198(2006).
Hu R.-M., et al. Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).
Bechtel S., et al. BMC Genomics 8:399-399(2007).
Higgins J.J., et al. Neurology 63:1927-1931(2004).