

**Pcsk1n Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14369**

**Specification**

---

**Pcsk1n Antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O9OXU9</a>
Other Accession	<a href="#">NM_019279</a> , <a href="#">NP_062152</a>
Reactivity	Human, Mouse, Rat, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28kDa kDa

**Pcsk1n Antibody - N-terminal region - Additional Information**

**Gene ID** 246333

**Alias Symbol** Saas

**Other Names**

ProSAAS, Proprotein convertase subtilisin/kexin type 1 inhibitor, Proprotein convertase 1 inhibitor, pro-SAAS, KEP, Big SAAS, b-SAAS, Little SAAS, I-SAAS, Big PEN-LEN, b-PEN-LEN, SAAS CT(1-49), PEN, PEN-20, Little LEN, I-LEN, Big LEN, b-LEN, SAAS CT(25-40), Pcsk1n

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-Pcsk1n antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

Pcsk1n Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**Pcsk1n Antibody - N-terminal region - Protein Information**

**Name** Pcsk1n

**Function**

May function in the control of the neuroendocrine secretory pathway. Proposed be a specific endogenous inhibitor of PCSK1 (PubMed: [10632593](http://www.uniprot.org/citations/10632593), PubMed: [10816562](http://www.uniprot.org/citations/10816562)). ProSAAS and Big PEN-LEN, both containing the C-terminal inhibitory domain, but not the processed peptides reduce PCSK1 activity in the endoplasmic

reticulum and Golgi. It reduces the activity of the 87 kDa form but not the autocatalytically derived 65 kDa form of PCSK1. Subsequent processing of proSAAS may eliminate the inhibition. Slows down convertase-mediated processing of proopiomelanocortin and proenkephalin. May control the intracellular timing of PCSK1 rather than its total level of activity (By similarity).

#### Cellular Location

Secreted {ECO:0000250|UniProtKB:Q9QXV0}. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:Q9QXV0}

#### Tissue Location


Expressed in adult brain (all major structural regions), adrenal gland (medulla) and spinal cord (dorsal and ventral horn). Expressed in pancreatic islands.

### Pcsk1n Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Pcsk1n Antibody - N-terminal region - Images



Host: Rabbit  
Target Name: Pcsk1n  
Sample Tissue: Rat Thymus lysates  
Antibody Dilution: 1.0µg/ml

### Pcsk1n Antibody - N-terminal region - References

Fricker L., et al. J. Neurosci. 20:639-648(2000).  
Donadel G., et al. Neuroendocrinology 67:190-196(1998).  
Qian Y., et al. J. Biol. Chem. 275:23596-23601(2000).  
Mzhavia N., et al. Biochem. J. 361:67-76(2002).  
Morgan D.J., et al. J. Neurochem. 93:1454-1462(2005).