

Brsk1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14473**Specification**

Brsk1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5RJI5
Other Accession	NM_001003920 , NP_001003920
Reactivity	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85kDa KDa

Brsk1 antibody - C-terminal region - Additional Information**Gene ID** 381979**Alias Symbol** Gm1100, MGC99905, SAD-B, SADB**Other Names**

Serine/threonine-protein kinase BRSK1, 2.7.11.1, 2.7.11.26, Brain-specific serine/threonine-protein kinase 1, BR serine/threonine-protein kinase 1, Serine/threonine-protein kinase SAD-B, Brsk1, Gm1100, Sadb

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Brsk1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Brsk1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Brsk1 antibody - C-terminal region - Protein Information**Name** Brsk1**Synonyms** Gm1100, Sadb**Function**

Serine/threonine-protein kinase that plays a key role in polarization of neurons and centrosome duplication. Phosphorylates CDC25B, CDC25C, MAPT/TAU, RIMS1, TUBG1, TUBG2 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of

cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-504' and 'Ser-554'. Also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in postmitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. In neurons, localizes to synaptic vesicles and plays a role in neurotransmitter release, possibly by phosphorylating RIMS1. Also acts as a positive regulator of centrosome duplication by mediating phosphorylation of gamma-tubulin (TUBG1 and TUBG2) at 'Ser-131', leading to translocation of gamma-tubulin and its associated proteins to the centrosome. Involved in the UV-induced DNA damage checkpoint response, probably by inhibiting CDK1 activity through phosphorylation and activation of WEE1, and inhibition of CDC25B and CDC25C.

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Synapse {ECO:0000250|UniProtKB:B2DD29}. Presynaptic active zone {ECO:0000250|UniProtKB:B2DD29}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:B2DD29} Note=Nuclear in the absence of DNA damage. Translocated to the nucleus in response to UV- or MMS-induced DNA damage (By similarity)

Tissue Location

Present in the gray matter of the brain and spinal cord (at protein level). Expressed in the nervous system, distributed within the brain and spinal cord of embryonic and postnatal animals

Brsk1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Brsk1 antibody - C-terminal region - Images



WB Suggested Anti-Brsk1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart

Brsk1 antibody - C-terminal region - References

Kishi M., et al. Science 307:929-932(2005).
Alvarado-Kristensson M., et al. Nat. Cell Biol. 11:1081-1092(2009).

Barnes A.P., et al. Cell 129:549-563(2007).
Muller M., et al. J. Cell Sci. 123:286-294(2010).