

FLT1 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14601**Specification**

FLT1 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P17948-6
Reactivity	Human, Rat, Rabbit, Goat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rat, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50kDa KDa

FLT1 Antibody - C-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-FLT1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

FLT1 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FLT1 Antibody - C-terminal region - Protein Information**FLT1 Antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FLT1 Antibody - C-terminal region - Images




168 kDa_
144 kDa_
90 kDa_
65 kDa_
40 kDa_

Host: Rabbit

Target Name: FLT1

Sample Tissue: HepG2 Whole cell lysate

s

Antibody Dilution: 1.0µg/ml There is BioGPS gene expression data showing that FLT1 is expressed in HepG2