

**VASP antibody - C-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14633**

**Specification**

---

**VASP antibody - C-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">P50552</a>
Other Accession	<a href="#">NM_003370</a> , <a href="#">NP_003361</a>
Reactivity	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

**VASP antibody - C-terminal region - Additional Information**

**Gene ID** 7408

**Other Names**

Vasodilator-stimulated phosphoprotein, VASP, VASP

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-VASP antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

VASP antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**VASP antibody - C-terminal region - Protein Information**

**Name** VASP

**Function**

Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance, lamellipodial and filopodial dynamics, platelet activation and cell migration. VASP promotes actin filament elongation. It protects the barbed end of growing actin filaments against capping and increases the rate of actin polymerization in the presence of capping protein. VASP stimulates actin filament elongation by promoting the transfer of profilin-bound actin monomers onto the barbed end of growing actin filaments. Plays a role in actin-based mobility of *Listeria monocytogenes* in host cells. Regulates actin dynamics in platelets and plays an important role in regulating platelet aggregation.

### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cell junction, focal adhesion. Cell junction, tight junction Cell projection, lamellipodium membrane. Cell projection, filopodium membrane. Note=Targeted to stress fibers and focal adhesions through interaction with a number of proteins including MRL family members Localizes to the plasma membrane in protruding lamellipodia and filopodial tips. Stimulation by thrombin or PMA, also translocates VASP to focal adhesions. Localized along the sides of actin filaments throughout the peripheral cytoplasm under basal conditions. In pre-apoptotic cells, colocalizes with MEFV in large specks (pyroptosomes)

### Tissue Location


Highly expressed in platelets.

### VASP antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### VASP antibody - C-terminal region - Images



WB Suggested Anti-VASP Antibody Titration: 1.0 µg/ml  
Positive Control: Jurkat Whole Cell

### VASP antibody - C-terminal region - References

- Haffner C.,et al.EMBO J. 14:19-27(1995).  
Laurent V.,et al.J. Cell Biol. 144:1245-1258(1999).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Gevaert K.,et al.Nat. Biotechnol. 21:566-569(2003).