

BNIP2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14736

Specification

BNIP2 Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q12982
Reactivity	Human
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34kDa KDa

BNIP2 Antibody - C-terminal region - Additional Information

Gene ID 663

Alias Symbol **BNIP2, NIP2,**
Other Names
BCL2/adenovirus E1B 19 kDa protein-interacting protein 2, BNIP2, NIP2

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 μ l of distilled water. Final Anti-BNIP2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

BNIP2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

BNIP2 Antibody - C-terminal region - Protein Information

Name BNIP2

Synonyms NIP2

Function

Implicated in the suppression of cell death. Interacts with the BCL-2 and adenovirus E1B 19 kDa proteins.

Cellular Location


Cytoplasm. Cytoplasm, perinuclear region. Note=Localizes to the nuclear envelope region and to other cytoplasmic structures

BNIP2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BNIP2 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: BNIP2
Sample Tissue: 721_B Whole Cell lysates
Antibody Dilution: 1.0µg/ml

BNIP2 Antibody - C-terminal region - References

- Boyd J.M., et al. Cell 79:341-351(1994).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Zody M.C., et al. Nature 440:671-675(2006).
Zahedi R.P., et al. J. Proteome Res. 7:526-534(2008).
Daub H., et al. Mol. Cell 31:438-448(2008).