

RPL37 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI14741**Specification**

RPL37 antibody - middle region - Product Information

Application	WB
Primary Accession	P61927
Other Accession	NM_000997 , NP_000988
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Chicken, Horse, Yeast, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11kDa KDa

RPL37 antibody - middle region - Additional Information**Gene ID 6167**

Alias Symbol **DKFZp686G1699, MGC99572, L37**
Other Names
60S ribosomal protein L37, G1.16, RPL37

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-RPL37 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

RPL37 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

RPL37 antibody - middle region - Protein Information**Name RPL37****Function**

Component of the large ribosomal subunit (PubMed: [23636399](http://www.uniprot.org/citations/23636399), PubMed: [32669547](http://www.uniprot.org/citations/32669547)). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: [23636399](http://www.uniprot.org/citations/23636399), PubMed: [32669547](http://www.uniprot.org/citations/32669547)).

Cellular Location


Cytoplasm.

RPL37 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RPL37 antibody - middle region - Images



WB Suggested Anti-RPL37 Antibody Titration: 1.0 μ g/ml

Positive Control: MCF7 Whole Cell

RPL37 antibody - middle region - References

- Kato S., et al. Submitted (NOV-1993) to the EMBL/GenBank/DDBJ databases.
Barnard G.F., et al. Biochim. Biophys. Acta 1218:425-428(1994).
Su S., et al. Eur. J. Biochem. 232:789-797(1995).
Yoshihama M., et al. Genome Res. 12:379-390(2002).
Ebert L., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.