

**RPA2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI14831****Specification**

---

**RPA2 antibody - N-terminal region - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O76021</a>
Other Accession	<a href="#">NM_015659</a> , <a href="#">NP_056474</a>
Reactivity	<b>Human, Mouse, Rat, Pig, Bovine, Guinea Pig, Dog</b>
Predicted Host	<b>Human, Mouse, Pig, Bovine</b>
Clonality	<b>Rabbit</b>
Calculated MW	<b>Polyclonal</b> <b>30kDa KDa</b>

**RPA2 antibody - N-terminal region - Additional Information****Gene ID** 26156**Alias Symbol** **REPA2, RPA32****Other Names**

Ribosomal L1 domain-containing protein 1, CATX-11, Cellular senescence-inhibited gene protein, Protein PBK1, RSL1D1

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-RPA2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

RPA2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**RPA2 antibody - N-terminal region - Protein Information****Name** RSL1D1**Synonyms** CATX11, CSIG {ECO:0000303|PubMed:1867864}**Function**

Regulates cellular senescence through inhibition of PTEN translation. Acts as a pro-apoptotic regulator in response to DNA damage.

**Cellular Location**

Nucleus, nucleolus. Note=Colocalizes with ING1 in the nucleolus after UV stress.


**Tissue Location**

Expressed at high intensities in the heart, skeletal muscle, and placenta.

**RPA2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**RPA2 antibody - N-terminal region - Images**

WB Suggested Anti-RPA2 Antibody Titration: 1.0 µg/ml  
Positive Control: Fetal Brain

**RPA2 antibody - N-terminal region - References**

- Huch G., et al. Placenta 19:557-567(1998).  
Petroziello J., et al. Oncogene 23:7734-7745(2004).  
Guo S.-Z., et al. Submitted (SEP-2002) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Martin J., et al. Nature 432:988-994(2004).