

CAMK2G antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14897

Specification

CAMK2G antibody - C-terminal region - Product Information

Application	WB
Primary Accession	O13555-8
Other Accession	NM_172171 , NP_751911
Reactivity	Human, Rat, Pig, Horse, Bovine, Dog
Predicted	Human, Rat, Pig, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61kDa KDa

CAMK2G antibody - C-terminal region - Additional Information

Alias Symbol	CAMK, CAMK-II, CAMKG, FLJ16043, MGC26678
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CAMK2G antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CAMK2G antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


CAMK2G antibody - C-terminal region - Protein Information

CAMK2G antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CAMK2G antibody - C-terminal region - Images



90 kDa
65 kDa
40 kDa
29 kDa
22 kDa

WB Suggested Anti-CAMK2G Antibody Titration: 1.0 μ g/ml
Positive Control: THP-1 Whole Cell