

CD58 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14932**Specification**

CD58 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P19256-3
Other Accession	NM_001144822 , NP_001138294
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27kDa KDa

CD58 antibody - C-terminal region - Additional Information

Alias Symbol **FLJ23181, FLJ43722, LFA-3, LFA3, ag3**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-CD58 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

CD58 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CD58 antibody - C-terminal region - Protein Information**CD58 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD58 antibody - C-terminal region - Images




WB Suggested Anti-CD58 Antibody Titration: 1.0 $\mu\text{g/ml}$

Positive Control: COLO205 Whole Cell CD58 is strongly supported by BioGPS gene expression data to be expressed in Human COLO205 cells