

MCAM Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI14968

Specification

MCAM Antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P43121-2
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Guinea Pig, Dog
Predicted Host	Human, Mouse, Rat, Rabbit, Pig, Horse
Clonality	Rabbit
Calculated MW	Polyclonal 57kDa KDa

MCAM Antibody - C-terminal region - Additional Information

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MCAM antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MCAM Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


MCAM Antibody - C-terminal region - Protein Information

MCAM Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MCAM Antibody - C-terminal region - Images





Host: Rabbit

Target Name: MCAM

Sample Tissue: HepG2 Whole cell lysate

S

Antibody Dilution: 1.0 μ g/ml There is BioGPS gene expression data showing that MCAM is expressed in HepG2