10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# FGF18 Antibody - C-terminal region
**Rabbit Polyclonal Antibody**
**Catalog # AI15026**

## Specification

### FGF18 Antibody - C-terminal region - Product Information

| | |
|---|---|
| Application | **WB** |
| Primary Accession | O76093 |
| Other Accession | NM_003862, NP_003853 |
| Reactivity | **Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog** |
| Predicted | **Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog** |
| Host | **Rabbit** |
| Clonality | **Polyclonal** |
| Calculated MW | **21kDa KDa** |

### FGF18 Antibody - C-terminal region - Additional Information

**Gene ID** 8817

| | |
|---|---|
| Alias Symbol | **FGF-18, ZFGF5** |

**Other Names**
Fibroblast growth factor 18, FGF-18, zFGF5, FGF18

**Format**
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**
Add 50 ul of distilled water. Final anti-FGF18 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**
FGF18 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

### FGF18 Antibody - C-terminal region - Protein Information

**Name** FGF18

**Function**
Plays an important role in the regulation of cell proliferation, cell differentiation and cell migration. Required for normal ossification and bone development. Stimulates hepatic and intestinal proliferation.

**Cellular Location**
Secreted.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

## FGF18 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## FGF18 Antibody - C-terminal region - Images




WB Suggested Anti-FGF18 Antibody Titration: 1.0 µg/ml
Positive Control: ACHN Whole Cell

## FGF18 Antibody - C-terminal region - References

Hu M.C.-T.,et al.Mol. Cell. Biol. 18:6063-6074(1998).
Ohbayashi N.,et al.J. Biol. Chem. 273:18161-18164(1998).
Deisher T.,et al.Submitted (DEC-1999) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.