

**FKBP10 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15187****Specification**

---

**FKBP10 antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O96AY3</a>
Other Accession	<a href="#">NM_021939</a> , <a href="#">NP_068758</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	61kDa kDa

**FKBP10 antibody - N-terminal region - Additional Information****Gene ID** 60681**Alias Symbol** FKBP65, FLJ20683, FLJ22041, FLJ23833, PPIASE, hFKBP65, OI6, OI11**Other Names**

Peptidyl-prolyl cis-trans isomerase FKBP10, PPIase FKBP10, 5.2.1.8, 65 kDa FK506-binding protein, 65 kDa FKBP, FKBP-65, FK506-binding protein 10, FKBP-10, Immunophilin FKBP65, Rotamase, FKBP10, FKBP65

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-FKBP10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

FKBP10 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**FKBP10 antibody - N-terminal region - Protein Information****Name** FKBP10**Synonyms** FKBP65**Function**

PPIases accelerate the folding of proteins during protein synthesis.

### Cellular Location


Endoplasmic reticulum lumen {ECO:0000255|PROSITE- ProRule:PRU10138}

### FKBP10 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### FKBP10 antibody - N-terminal region - Images



WB Suggested Anti-FKBP10 Antibody Titration: 1.0  $\mu$ g/ml  
Positive Control: Placenta

### FKBP10 antibody - N-terminal region - References

- Rulten S., et al. Submitted (JAN-2001) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Otsuki T., et al. DNA Res. 12:117-126(2005).  
Bechtel S., et al. BMC Genomics 8:399-399(2007).  
Zody M.C., et al. Nature 440:1045-1049(2006).