

**BBS9 Antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI15282****Specification**

---

**BBS9 Antibody - N-terminal region - Product Information**

Application	WB
Primary Accession	<a href="#">O3SYG4</a>
Other Accession	<a href="#">NM_014451</a> , <a href="#">NP_055266</a>
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	95kDa KDa

**BBS9 Antibody - N-terminal region - Additional Information****Gene ID** 27241**Alias Symbol** **B1, C18, D1, MGC118917, PTHB1**  
**Other Names**  
Protein PTHB1, Bardet-Biedl syndrome 9 protein, Parathyroid hormone-responsive B1 gene protein, BBS9, PTHB1**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-BBS9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

BBS9 Antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**BBS9 Antibody - N-terminal region - Protein Information****Name** BBS9**Synonyms** PTHB1**Function**

The BBSome complex is thought to function as a coat complex required for sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome. Rab8(GTP)

enters the primary cilium and promotes extension of the ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to RAB31P/Rabin8, the guanosyl exchange factor (GEF) for Rab8 and then the Rab8-GTP localizes to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary membrane. Required for proper BBSome complex assembly and its ciliary localization.

#### Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium membrane. Cytoplasm Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite

#### Tissue Location


Widely expressed. Expressed in adult heart, skeletal muscle, lung, liver, kidney, placenta and brain, and in fetal kidney, lung, liver and brain.

### BBS9 Antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### BBS9 Antibody - N-terminal region - Images



WB Suggested Anti-BBS9 Antibody Titration: 1.0 µg/ml  
Positive Control: Jurkat Whole Cell

### BBS9 Antibody - N-terminal region - References

- Hillier L.W., et al. Nature 424:157-164(2003).  
Adams A.E., et al. Bone 24:305-313(1999).  
Keen T.J., et al. Submitted (JAN-1997) to the EMBL/GenBank/DDBJ databases.  
Vernon E.G., et al. Oncogene 22:1371-1380(2003).  
Nachury M.V., et al. Cell 129:1201-1213(2007).