10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FSD2 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI15362

Specification

FSD2 Antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	A1L4K1
Other Accession	NM_001007122, NP_001007123
Reactivity	**Human, Rat, Pig, Horse, Bovine, Guinea Pig, Dog**
Predicted	**Human, Rat, Pig, Horse, Bovine, Guinea Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**85kDa KDa**

FSD2 Antibody - C-terminal region - Additional Information

Gene ID 123722

Alias Symbol	**RP11-127F21, SPRYD1**

Other Names
Fibronectin type III and SPRY domain-containing protein 2, SPRY domain-containing protein 1, FSD2, SPRYD1

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-FSD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
FSD2 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FSD2 Antibody - C-terminal region - Protein Information

Name FSD2

Synonyms SPRYD1

Cellular Location
Nucleus {ECO:0000250|UniProtKB:H0UZ81}. Sarcoplasmic reticulum {ECO:0000250|UniProtKB:H0UZ81}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:H0UZ81}. Note=In skeletal muscles and striated muscles flanks Z-disks. Partially colocalizes with RYR2 in the sarcoplasmic reticulum. {ECO:0000250|UniProtKB:H0UZ81}

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FSD2 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FSD2 Antibody - C-terminal region - Images



WB Suggested Anti-FSD2 Antibody Titration: 1.0 µg/ml
Positive Control: 721_B Whole Cell

FSD2 Antibody - C-terminal region - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Zody M.C.,et al.Nature 440:671-675(2006).