

IGF2BP3 Antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI16033**Specification****IGF2BP3 Antibody - middle region - Product Information**

Application	WB
Primary Accession	O00425
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63kDa KDa

IGF2BP3 Antibody - middle region - Additional Information**Gene ID** 10643**Alias Symbol** **IGF2BP3, IMP3, KOC1, VICKZ3,**
Other Names

Insulin-like growth factor 2 mRNA-binding protein 3, IGF2 mRNA-binding protein 3, IMP-3, IGF-II mRNA-binding protein 3, KH domain-containing protein overexpressed in cancer, hKOC, VICKZ family member 3, IGF2BP3, IMP3, KOC1, VICKZ3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 µl of distilled water. Final Anti-IGF2BP3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.

Precautions

IGF2BP3 Antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

IGF2BP3 Antibody - middle region - Protein Information**Name** IGF2BP3**Synonyms** IMP3, KOC1, VICKZ3**Function**

RNA-binding factor that may recruit target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation. Preferentially binds to N6- methyladenosine (m6A)-containing mRNAs and increases their stability (PubMed:29476152). Binds to the

3'-UTR of CD44 mRNA and stabilizes it, hence promotes cell adhesion and invadopodia formation in cancer cells. Binds to beta-actin/ACTB and MYC transcripts. Increases MYC mRNA stability by binding to the coding region instability determinant (CRD) and binding is enhanced by m6A-modification of the CRD (PubMed: <http://www.uniprot.org/citations/29476152> target="_blank">29476152). Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs.

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, P-body. Cytoplasm, Stress granule. Note=Found in lamellipodia of the leading edge, in the perinuclear region, and beneath the plasma membrane. The subcytoplasmic localization is cell specific and regulated by cell contact and growth. Localized at the connecting piece and the tail of the spermatozoa. Colocalized with CD44 mRNA in RNP granules. In response to cellular stress, such as oxidative stress, recruited to stress granules

Tissue Location


Expressed in fetal liver, fetal lung, fetal kidney, fetal thymus, fetal placenta, fetal follicles of ovary and gonocytes of testis, growing oocytes, spermatogonia and semen (at protein level) Expressed in cervix adenocarcinoma, in testicular, pancreatic and renal-cell carcinomas (at protein level). Expressed ubiquitously during fetal development at 8 and 14 weeks of gestation. Expressed in ovary, testis, brain, placenta, pancreatic cancer tissues and pancreatic cancer cell lines.

IGF2BP3 Antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IGF2BP3 Antibody - middle region - Images



Host: Rabbit
Target Name: IGF2BP3
Sample Tissue: 721_B Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

IGF2BP3 Antibody - middle region - Background

RNA-binding factor that may recruits target transcripts to cytoplasmic protein-RNA complexes (mRNPs). This transcript 'caging' into mRNPs allows mRNA transport and transient storage. It also modulates the rate and location at which target transcripts encounter the translational apparatus and shields them from endonuclease attacks or microRNA-mediated degradation. Binds to the 3'-UTR of CD44 mRNA and stabilizes it, hence promotes cell adhesion and invadopodia formation in cancer cells. Binds to beta- actin/ACTB and MYC transcripts. Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs.

IGF2BP3 Antibody - middle region - References

Mueller-Pillasch F.,et al.Oncogene 14:2729-2733(1997).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Hillier L.W.,et al.Nature 424:157-164(2003).
Mueller-Pillasch F.,et al.Mech. Dev. 88:95-99(1999).
Nielsen J.,et al.Mol. Cell. Biol. 19:1262-1270(1999).