

CDK10 Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI16062**Specification**

CDK10 Antibody - C-terminal region - Product Information

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | Q15131 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 39kDa KDa |

CDK10 Antibody - C-terminal region - Additional Information**Gene ID** 8558**Alias Symbol** CDK10,
Other Names

Cyclin-dependent kinase 10, 2.7.11.22, Cell division protein kinase 10, Serine/threonine-protein kinase PISSLRE, CDK10

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & StorageAdd 50 μ l of distilled water. Final Anti-CDK10 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at -20°C. Avoid repeat freeze-thaw cycles.**Precautions**

CDK10 Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CDK10 Antibody - C-terminal region - Protein Information**Name** CDK10**Function**Cyclin-dependent kinase that phosphorylates the transcription factor ETS2 (in vitro) and positively controls its proteasomal degradation (in cells) (PubMed:[24218572](http://www.uniprot.org/citations/24218572)). Involved in the regulation of actin cytoskeleton organization through the phosphorylation of actin dynamics regulators such as PKN2. Is a negative regulator of ciliogenesis through phosphorylation of PKN2 and promotion of RhoA signaling (PubMed:[27104747](http://www.uniprot.org/citations/27104747)).**Cellular Location**


Cytoplasm, cytoskeleton, cilium basal body

CDK10 Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDK10 Antibody - C-terminal region - Images



Host: Rabbit
Target Name: CDK10
Sample Tissue: MCF7 Whole cell lysate
s
Antibody Dilution: 1.0µg/ml

CDK10 Antibody - C-terminal region - Background

Cyclin-dependent kinase that phosphorylates the transcription factor ETS2 (in vitro) and positively controls its proteasomal degradation (in cells).

CDK10 Antibody - C-terminal region - References

- Grana X.,et al.Oncogene 9:2097-2103(1994).
Grana X.,et al.Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
Brambilla R.,et al.Oncogene 9:3037-3041(1994).
Crawford J.,et al.Genomics 56:90-97(1999).
Ota T.,et al.Nat. Genet. 36:40-45(2004).