

**CNR2 / CB2 Antibody (Internal)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10013**

**Specification**

---

**CNR2 / CB2 Antibody (Internal) - Product Information**

Application	IHC
Primary Accession	<a href="#">P34972</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40kDa KDa

**CNR2 / CB2 Antibody (Internal) - Additional Information**

**Gene ID** 1269

**Other Names**

Cannabinoid receptor 2, CB-2, CB2, hCB2, CX5, CNR2, CB2A, CB2B

**Target/Specificity**

Human CNR2 / CB2. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except PROS1 (47%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

CNR2 / CB2 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

**CNR2 / CB2 Antibody (Internal) - Protein Information**

**Name** CNR2

**Synonyms** CB2A, CB2B

**Function**

Heterotrimeric G protein-coupled receptor for endocannabinoid 2-arachidonoylglycerol mediating inhibition of adenylate cyclase. May function in inflammatory response, nociceptive transmission and bone homeostasis.

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite. Perikaryon Note=Localizes to apical dendrite of pyramidal neurons.

**Tissue Location**

Preferentially expressed in cells of the immune system with higher expression in B-cells and NK

cells (at protein level). Expressed in skin in suprabasal layers and hair follicles (at protein level). Highly expressed in tonsil and to a lower extent in spleen, peripheral blood mononuclear cells, and thymus. PubMed:14657172 could not detect expression in normal brain. Expressed in brain by perivascular microglial cells and dorsal root ganglion sensory neurons (at protein level). Two isoforms are produced by alternative promoter usage and differ only in the 5' UTR: isoform CB2A is observed predominantly in testis with some expression in brain, while isoform CB2B is predominant in spleen and leukocytes


**Volume**  
50  $\mu$ l

### **CNR2 / CB2 Antibody (Internal) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### **CNR2 / CB2 Antibody (Internal) - Images**



Anti-CNR2 / CB2 antibody IHC of human tonsil.



Anti-CNR2 / CB2 antibody ALS10013 IHC of human germinal center and mantle zone.



Anti-CNR2 / CB2 antibody IHC of human Brain, Glioblastoma.

### **CNR2 / CB2 Antibody (Internal) - Background**

Heterotrimeric G protein-coupled receptor for endocannabinoid 2-arachidonoylglycerol mediating inhibition of adenylate cyclase. May function in inflammatory response, nociceptive transmission and bone homeostasis.

### **CNR2 / CB2 Antibody (Internal) - References**

- Munro S.,et al.Nature 365:61-65(1993).
- Liu Q.-R.,et al.Genes Brain Behav. 8:519-530(2009).
- Bruess M.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.
- Warren C.N.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.
- Saravanan T.,et al.Submitted (NOV-2005) to the EMBL/GenBank/DDBJ databases.