

**GRM1 / MGLUR1 Antibody (N-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10194****Specification**

---

**GRM1 / MGLUR1 Antibody (N-Terminus) - Product Information**

Application	IHC
Primary Accession	<a href="#">O13255</a>
Reactivity	Human, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	132kDa KDa

**GRM1 / MGLUR1 Antibody (N-Terminus) - Additional Information****Gene ID** 2911**Other Names**

Metabotropic glutamate receptor 1, mGluR1, GRM1, GPRC1A, MGLUR1

**Target/Specificity**

Human GRM1 / MGLUR1. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

GRM1 / MGLUR1 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**GRM1 / MGLUR1 Antibody (N-Terminus) - Protein Information****Name** GRM1**Synonyms** GPRC1A, MGLUR1**Function**

G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Signaling activates a phosphatidylinositol- calcium second messenger system. May participate in the central action of glutamate in the CNS, such as long-term potentiation in the hippocampus and long-term depression in the cerebellum (PubMed:<a href="http://www.uniprot.org/citations/24603153" target="\_blank">24603153</a>, PubMed:<a href="http://www.uniprot.org/citations/28886343" target="\_blank">28886343</a>, PubMed:<a href="http://www.uniprot.org/citations/7476890" target="\_blank">7476890</a>). May function in the light response in the retina (By similarity). Induces GRID1 and GRID2 cation-channel activation via GNAQ-PLC-PKC pathway in dopaminergic neurons and cerebellar Purkinje cell, respectively

(PubMed:<a href="http://www.uniprot.org/citations/24357660" target="\_blank">24357660</a>,  
PubMed:<a href="http://www.uniprot.org/citations/27276689" target="\_blank">27276689</a>).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane; Multi-pass membrane protein

#### Tissue Location

Detected in brain..

#### Volume


50  $\mu$ l

### GRM1 / MGLUR1 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GRM1 / MGLUR1 Antibody (N-Terminus) - Images



Anti-GRM1 / MGLUR1 antibody ALS10194 IHC of human brain, neurons and glia.

### GRM1 / MGLUR1 Antibody (N-Terminus) - Background

G-protein coupled receptor for glutamate. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Signaling activates a phosphatidylinositol-calcium second messenger system. May participate in the central action of glutamate in the CNS, such as long-term potentiation in the hippocampus and long-term depression in the cerebellum.

### GRM1 / MGLUR1 Antibody (N-Terminus) - References

Desai M.A., et al. Mol. Pharmacol. 48:648-657(1995).

Stephan D., et al. *Neuropharmacology* 35:1649-1660(1996).  
Mungall A.J., et al. *Nature* 425:805-811(2003).  
Guergueltcheva V., et al. *Am. J. Hum. Genet.* 91:553-564(2012).  
Wu H., et al. *Science* 344:58-64(2014).