

CCR3 Antibody (Extracellular Domain)
Rabbit Polyclonal Antibody
Catalog # ALS10347**Specification**

CCR3 Antibody (Extracellular Domain) - Product Information

Application	IHC
Primary Accession	P51677
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

CCR3 Antibody (Extracellular Domain) - Additional Information**Gene ID** 1232**Other Names**

C-C chemokine receptor type 3, C-C CKR-3, CC-CKR-3, CCR-3, CCR3, CKR3, Eosinophil eotaxin receptor, CD193, CCR3, CMKBR3

Target/Specificity

Human CCR3. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Long term: -70°C; Short term: +4°C

Precautions

CCR3 Antibody (Extracellular Domain) is for research use only and not for use in diagnostic or therapeutic procedures.

CCR3 Antibody (Extracellular Domain) - Protein Information**Name** CCR3**Synonyms** CMKBR3**Function**

Receptor for C-C type chemokine. Binds and responds to a variety of chemokines, including CCL11, CCL26, CCL7, CCL13, RANTES(CCL5) and CCL15 (PubMed:7622448, PubMed:8642344, PubMed:8676064). Subsequently transduces a signal by increasing the intracellular calcium ions level (PubMed:8676064). In addition acts as a possible functional receptor for NARS1 (PubMed:30171954).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

In eosinophils as well as trace amounts in neutrophils and monocytes.

Volume


50 μ l

CCR3 Antibody (Extracellular Domain) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CCR3 Antibody (Extracellular Domain) - Images



Anti-CCR3 antibody ALS10347 IHC of human nasal mucosa, allergic rhinitis, eosinophils.



Anti-CCR3 antibody IHC of human Pancreas, Carcinoma.



Anti-CCR3 antibody IHC of human tonsil.

CCR3 Antibody (Extracellular Domain) - Background

Receptor for a C-C type chemokine. Binds to eotaxin, eotaxin-3, MCP-3, MCP-4, RANTES and MIP-1 delta. Subsequently transduces a signal by increasing the intracellular calcium ions level. Alternative coreceptor with CD4 for HIV-1 infection.

CCR3 Antibody (Extracellular Domain) - References

Combadiere C., et al. J. Biol. Chem. 270:16491-16494(1995).
Combadiere C., et al. J. Biol. Chem. 270:30235-30235(1995).
Daugherty B.L., et al. J. Exp. Med. 183:2349-2354(1996).
Ponath P.D., et al. J. Exp. Med. 183:2437-2448(1996).
Xiao L., et al. Submitted (NOV-1997) to the EMBL/GenBank/DDBJ databases.