

**NR4A1 / NUR77 Antibody (N-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10653****Specification**

---

**NR4A1 / NUR77 Antibody (N-Terminus) - Product Information**

Application	IHC
Primary Accession	<a href="#">P22736</a>
Reactivity	Human, Mouse, Hamster
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64kDa KDa

**NR4A1 / NUR77 Antibody (N-Terminus) - Additional Information****Gene ID** 3164**Other Names**

Nuclear receptor subfamily 4 group A member 1, Early response protein NAK1, Nuclear hormone receptor NUR/77, Nur77, Orphan nuclear receptor HMR, Orphan nuclear receptor TR3, ST-59, Testicular receptor 3, NR4A1, GFRP1, HMR, NAK1

**Target/Specificity**

Human NUR77. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

NR4A1 / NUR77 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**NR4A1 / NUR77 Antibody (N-Terminus) - Protein Information****Name** NR4A1**Synonyms** GFRP1, HMR, NAK1**Function**

Orphan nuclear receptor. Binds the NGFI-B response element (NBRE) 5'-AAAGGTCA-3' (PubMed:[18690216](http://www.uniprot.org/citations/18690216)), PubMed:[8121493](http://www.uniprot.org/citations/8121493), PubMed:[9315652](http://www.uniprot.org/citations/9315652)). Binds 9-cis-retinoic acid outside of its ligand-binding (NR LBD) domain (PubMed:[18690216](http://www.uniprot.org/citations/18690216)). Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation (PubMed:[22983157](http://www.uniprot.org/citations/22983157))

target="\_blank">22983157</a>). Regulates the inflammatory response in macrophages by regulating metabolic adaptations during inflammation, including repressing the transcription of genes involved in the citric acid cycle (TCA) (By similarity). Inhibits NF-kappa-B signaling by binding to low-affinity NF-kappa-B binding sites, such as at the IL2 promoter (PubMed:<a href="http://www.uniprot.org/citations/15466594" target="\_blank">15466594</a>). May act concomitantly with NR4A2 in regulating the expression of delayed-early genes during liver regeneration (By similarity). Plays a role in the vascular response to injury (By similarity).

#### Cellular Location

Nucleus. Cytoplasm, cytosol. Mitochondrion Note=Nuclear export to the cytosol is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340). Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950).

#### Tissue Location

Fetal muscle and adult liver, brain and thyroid.

#### Volume


50 µl

### NR4A1 / NUR77 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NR4A1 / NUR77 Antibody (N-Terminus) - Images



Anti-NUR77 antibody ALS10653 IHC of human ovary.

### NR4A1 / NUR77 Antibody (N-Terminus) - Background

Orphan nuclear receptor. May act concomitantly with NURR1 in regulating the expression of delayed-early genes during liver regeneration. Binds the NGFI-B response element (NBRE) 5'-AAAAGGTCA-3' (By similarity). May inhibit NF-kappa-B transactivation of IL2. Participates in energy

homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation.

#### **NR4A1 / NUR77 Antibody (N-Terminus) - References**

Nakai A., et al. Mol. Endocrinol. 4:1438-1443(1990).

Chang C., et al. J. Steroid Biochem. 34:391-395(1989).

Ohkura N., et al. Submitted (MAY-1996) to the EMBL/GenBank/DDBJ databases.

Kobayashi T., et al. Submitted (DEC-2008) to the EMBL/GenBank/DDBJ databases.

Kaighin V.A., et al. Submitted (DEC-2010) to the EMBL/GenBank/DDBJ databases.