

**ADGRG1 / GPR56 Antibody (N-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS10723****Specification****ADGRG1 / GPR56 Antibody (N-Terminus) - Product Information**

Application	IHC
Primary Accession	<a href="#">O9Y653</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa

**ADGRG1 / GPR56 Antibody (N-Terminus) - Additional Information****Gene ID** 9289**Other Names**

G-protein coupled receptor 56, Protein TM7XN1, GPR56 N-terminal fragment, GPR56 NT, GPR56(N), GPR56 extracellular subunit, GPR56 subunit alpha, GPR56 C-terminal fragment, GPR56 CT, GPR56(C), GPR56 seven-transmembrane subunit, GPR56 7TM, GPR56 subunit beta, GPR56, TM7LN4, TM7XN1

**Target/Specificity**

Human GPR56. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except MACF1 (44%).

**Reconstitution & Storage**

Long term: -70°C; Short term: +4°C

**Precautions**

ADGRG1 / GPR56 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**ADGRG1 / GPR56 Antibody (N-Terminus) - Protein Information****Name** ADGRG1 ([HGNC:4512](#))**Synonyms** GPR56, TM7LN4, TM7XN1**Function**

Receptor involved in cell adhesion and probably in cell-cell interactions. Mediates cell matrix adhesion in developing neurons and hematopoietic stem cells. Receptor for collagen III/COL3A1 in the developing brain and involved in regulation of cortical development, specifically in maintenance of the pial basement membrane integrity and in cortical lamination (By similarity). Binding to the COL3A1 ligand inhibits neuronal migration and activates the RhoA pathway by coupling to GNA13 and possibly GNA12 (PubMed:<a href="http://www.uniprot.org/citations/22238662" target="\_blank">22238662</a>). Plays a role

in the maintenance of hematopoietic stem cells and/or leukemia stem cells in bone marrow niche (By similarity). Plays a critical role in cancer progression by inhibiting VEGFA production thereby inhibiting angiogenesis through a signaling pathway mediated by PRKCA (PubMed:<a href="http://www.uniprot.org/citations/16757564" target="\_blank">16757564</a>, PubMed:<a href="http://www.uniprot.org/citations/21724588" target="\_blank">21724588</a>). Plays an essential role in testis development (By similarity).

#### Cellular Location

Cell membrane; Multi-pass membrane protein [ADGRG1 C-terminal fragment]: Membrane raft. Note=Interaction with its ligand COL3A1 leads to the release of ADGRG1 NT from the membrane and triggers the association of ADGRG1 CT with lipid rafts

#### Tissue Location

Widely distributed with highest levels found in thyroid gland, brain and heart. Expressed in a great number of tumor cells. Expression is down-regulated in different tumors from highly metastatic cells.

#### Volume


50 µl

### ADGRG1 / GPR56 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ADGRG1 / GPR56 Antibody (N-Terminus) - Images



Human, Prostate: Formalin-Fixed Paraffin-Embedded (FFPE)

### ADGRG1 / GPR56 Antibody (N-Terminus) - Background

Involved in cell adhesion and probably in cell-cell interactions. Regulates the migration of neural precursor cells. Receptor for collagen III/COL3A1 in the developing brain and involved in regulation of cortical development, specifically in maintenance of the pial basement membrane integrity and

in cortical lamination. Binding to the COL3A1 ligand inhibits neuronal migration and activates the RhoA pathway by coupling to GNA13 and possibly GNA12. Isoforms show differences in receptor signaling, specifically in serum response element (SRE) transcriptional activation upon overexpression. Overexpression inhibits melanoma tumor growth and metastasis and, during melanoma progression, regulates VEGFA production and angiogenesis through PRKCA; unprocessed GPR56 is inhibiting and GPR56 NT is activating angiogenesis. Required for normal cortical development and regulation of neuroprogenitor cells proliferation.

#### **ADGRG1 / GPR56 Antibody (N-Terminus) - References**

- Liu M., et al. Genomics 55:296-305(1999).  
Zendman A.J.W., et al. FEBS Lett. 446:292-298(1999).  
Kaighin V.A., et al. Submitted (DEC-2007) to the EMBL/GenBank/DDBJ databases.  
Clark H.F., et al. Genome Res. 13:2265-2270(2003).  
Ota T., et al. Nat. Genet. 36:40-45(2004).