

CARMA1 / CARD11 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS11402**Specification****CARMA1 / CARD11 Antibody (C-Terminus) - Product Information**

Application	IF, WB
Primary Accession	O9BXL7
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	133kDa KDa

CARMA1 / CARD11 Antibody (C-Terminus) - Additional Information**Gene ID** 84433**Other Names**

Caspase recruitment domain-containing protein 11, CARD-containing MAGUK protein 1, Carma 1, CARD11, CARMA1

Target/Specificity

synthetic peptide corresponding to amino acids at the C-terminus of human CARMA1

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

CARMA1 / CARD11 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CARMA1 / CARD11 Antibody (C-Terminus) - Protein Information**Name** CARD11 {ECO:0000303|PubMed:11278692, ECO:0000312|HGNC:HGNC:16393}**Function**

Adapter protein that plays a key role in adaptive immune response by transducing the activation of NF-kappa-B downstream of T- cell receptor (TCR) and B-cell receptor (BCR) engagement (PubMed:11278692, PubMed:11356195, PubMed:12356734). Transduces signals downstream TCR or BCR activation via the formation of a multiprotein complex together with BCL10 and MALT1 that induces NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways (PubMed:11356195). Upon activation in response to TCR or BCR triggering, CARD11 homooligomerizes to form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10 and subsequent recruitment of MALT1: this leads to I-kappa-B kinase (IKK) phosphorylation and degradation, and release of NF-kappa-B

proteins for nuclear translocation (PubMed:24074955). Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner (PubMed:17287217). Promotes linear ubiquitination of BCL10 by promoting the targeting of BCL10 to RNF31/HOIP (PubMed:27777308). Stimulates the phosphorylation of BCL10 (PubMed:11356195). Also activates the TORC1 signaling pathway (PubMed:28628108).

Cellular Location

Cytoplasm. Membrane raft. Note=Colocalized with DPP4 in membrane rafts.

Tissue Location


Detected in adult peripheral blood leukocytes, thymus, spleen and liver. Also found in promyelocytic leukemia HL-60 cells, chronic myelogenous leukemia K-562 cells, Burkitt's lymphoma Raji cells and colorectal adenocarcinoma SW480 cells. Not detected in HeLaS3, MOLT-4, A-549 and G431 cells.

CARMA1 / CARD11 Antibody (C-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CARMA1 / CARD11 Antibody (C-Terminus) - Images



Immunofluorescence of CARMA1 in Mouse Spleen cells with CARMA1 antibody at 20 ug/ml.



Western blot of CARMA1 expression in mouse thymus cell lysate with CARMA1 antibody at 0.5 (lane...

CARMA1 / CARD11 Antibody (C-Terminus) - Background

Involved in the costimulatory signal essential for T- cell receptor (TCR)-mediated T-cell activation. Its binding to DPP4 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Activates NF-kappa-B via BCL10 and IKK. Stimulates the phosphorylation of BCL10.

CARMA1 / CARD11 Antibody (C-Terminus) - References

- Bertin J.,et al.J. Biol. Chem. 276:11877-11882(2001).
- Hillier L.W.,et al.Nature 424:157-164(2003).
- Scherer S.W.,et al.Science 300:767-772(2003).
- Gaide O.,et al.FEBS Lett. 496:121-127(2001).
- Gaide O.,et al.FEBS Lett. 505:198-198(2001).