

**PDCD1 / CD279 / PD-1 Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS11431**

### Specification

---

#### PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Product Information

Application	IF
Primary Accession	<a href="#">Q15116</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

#### PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Additional Information

Gene ID 5133

#### Other Names

Programmed cell death protein 1, Protein PD-1, hPD-1, CD279, PDCD1, PD1

#### Target/Specificity

16 amino acid peptide from near the carboxy terminus of human PDCD1.

#### Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

#### Precautions

PDCD1 / CD279 / PD-1 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

#### PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Protein Information

**Name** PDCD1 {ECO:0000303|PubMed:7851902, ECO:0000312|HGNC:HGNC:8760}

#### Function

Inhibitory receptor on antigen activated T-cells that plays a critical role in induction and maintenance of immune tolerance to self (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>). Delivers inhibitory signals upon binding to ligands CD274/PDCD1L1 and CD273/PDCD1LG2 (PubMed:<a href="http://www.uniprot.org/citations/21276005" target="\_blank">21276005</a>). Following T-cell receptor (TCR) engagement, PDCD1 associates with CD3-TCR in the immunological synapse and directly inhibits T-cell activation (By similarity). Suppresses T-cell activation through the recruitment of PTPN11/SHP-2: following ligand-binding, PDCD1 is phosphorylated within the ITSM motif, leading to the recruitment of the protein tyrosine phosphatase PTPN11/SHP-2 that mediates dephosphorylation of key TCR proximal signaling molecules, such as ZAP70, PRKCQ/PKCtheta and CD247/CD3zeta (By similarity).

#### Cellular Location


Cell membrane; Single-pass type I membrane protein

### **PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Images**



Immunofluorescence of PD-1 in Human Brain cells with PD-1 antibody at 20 ug/ml.

### **PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - Background**

Inhibitory cell surface receptor involved in the regulation of T-cell function during immunity and tolerance. Upon ligand binding, inhibits T-cell effector functions in an antigen-specific manner. Possible cell death inducer, in association with other factors.

### **PDCD1 / CD279 / PD-1 Antibody (C-Terminus) - References**

Shinohara T., et al. *Genomics* 23:704-706(1994).  
Finger L.R., et al. *Gene* 197:177-187(1997).  
Finger L.R., et al. *Gene* 203:253-253(1997).  
Prokunina L., et al. *Nat. Genet.* 32:666-669(2002).  
He X., et al. Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.