

**BCL10 / BCL-10 Antibody (aa5-19)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS11548****Specification****BCL10 / BCL-10 Antibody (aa5-19) - Product Information**

Application	IHC
Primary Accession	<a href="#">O95999</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26kDa KDa

**BCL10 / BCL-10 Antibody (aa5-19) - Additional Information****Gene ID** 8915**Other Names**

B-cell lymphoma/leukemia 10, B-cell CLL/lymphoma 10, Bcl-10, CARD-containing molecule enhancing NF-kappa-B, CARD-like apoptotic protein, hCLAP, CED-3/ICH-1 prodomain homologous E10-like regulator, CIPER, Cellular homolog of vCARMEN, cCARMEN, Cellular-E10, c-E10, Mammalian CARD-containing adapter molecule E10, mE10, BCL10, CIPER, CLAP

**Target/Specificity**

Amino acids 5-19 of human BCL10

**Reconstitution & Storage**

+4°C or -20°C, Avoid repeated freezing and thawing.

**Precautions**

BCL10 / BCL-10 Antibody (aa5-19) is for research use only and not for use in diagnostic or therapeutic procedures.

**BCL10 / BCL-10 Antibody (aa5-19) - Protein Information****Name** BCL10 {ECO:0000303|PubMed:9989495, ECO:0000312|HGNC:HGNC:989}**Function**

Plays a key role in both adaptive and innate immune signaling by bridging CARD domain-containing proteins to immune activation (PubMed: <a href="http://www.uniprot.org/citations/10187770" target="\_blank">10187770</a>, PubMed: <a href="http://www.uniprot.org/citations/10364242" target="\_blank">10364242</a>, PubMed: <a href="http://www.uniprot.org/citations/10400625" target="\_blank">10400625</a>, PubMed: <a href="http://www.uniprot.org/citations/24074955" target="\_blank">24074955</a>, PubMed: <a href="http://www.uniprot.org/citations/25365219" target="\_blank">25365219</a>). Acts by channeling adaptive and innate immune signaling downstream of CARD domain-containing proteins CARD9, CARD11 and CARD14 to activate NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding

pro-inflammatory cytokines and chemokines (PubMed:<a href="http://www.uniprot.org/citations/24074955" target="\_blank">24074955</a>). Recruited by activated CARD domain-containing proteins: homooligomerized CARD domain-containing proteins form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10, subsequent recruitment of MALT1 and formation of a CBM complex (PubMed:<a href="http://www.uniprot.org/citations/24074955" target="\_blank">24074955</a>). This leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:<a href="http://www.uniprot.org/citations/18287044" target="\_blank">18287044</a>, PubMed:<a href="http://www.uniprot.org/citations/24074955" target="\_blank">24074955</a>, PubMed:<a href="http://www.uniprot.org/citations/27777308" target="\_blank">27777308</a>). Activated by CARD9 downstream of C-type lectin receptors; CARD9-mediated signals are essential for antifungal immunity (PubMed:<a href="http://www.uniprot.org/citations/26488816" target="\_blank">26488816</a>). Activated by CARD11 downstream of T-cell receptor (TCR) and B-cell receptor (BCR) (PubMed:<a href="http://www.uniprot.org/citations/18264101" target="\_blank">18264101</a>, PubMed:<a href="http://www.uniprot.org/citations/18287044" target="\_blank">18287044</a>, PubMed:<a href="http://www.uniprot.org/citations/24074955" target="\_blank">24074955</a>, PubMed:<a href="http://www.uniprot.org/citations/27777308" target="\_blank">27777308</a>). Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK (PubMed:<a href="http://www.uniprot.org/citations/10187815" target="\_blank">10187815</a>).

#### Cellular Location

Cytoplasm, perinuclear region. Membrane raft. Note=Appears to have a perinuclear, compact and filamentous pattern of expression. Also found in the nucleus of several types of tumor cells. Colocalized with DPP4 in membrane rafts.

#### Tissue Location

Ubiquitous..


#### BCL10 / BCL-10 Antibody (aa5-19) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### BCL10 / BCL-10 Antibody (aa5-19) - Images





Anti-BCL10 antibody IHC of human skeletal muscle.



Anti-BCL10 antibody IHC of human brain, cortex.

### **BCL10 / BCL-10 Antibody (aa5-19) - Background**

Involved in adaptive immune response (PubMed:25365219). Promotes apoptosis, pro-caspase-9 maturation and activation of NF- kappa-B via NIK and IKK. May be an adapter protein between upstream TNFR1-TRADD-RIP complex and the downstream NIK-IKK-IKAP complex. Is a substrate for MALT1 (PubMed:18264101).

### **BCL10 / BCL-10 Antibody (aa5-19) - References**

- Willis T.G.,et al.Cell 96:35-45(1999).
- Koseki T.,et al.J. Biol. Chem. 274:9955-9961(1999).
- Thome M.,et al.J. Biol. Chem. 274:9962-9968(1999).
- Yan M.,et al.J. Biol. Chem. 274:10287-10292(1999).
- Srinivasula S.M.,et al.J. Biol. Chem. 274:17946-17954(1999).