

Carboxylesterase 1 / CES1 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS12250**Specification****Carboxylesterase 1 / CES1 Antibody (Internal) - Product Information**

Application	WB, IHC
Primary Accession	P23141
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	63kDa KDa

Carboxylesterase 1 / CES1 Antibody (Internal) - Additional Information**Gene ID** 1066**Other Names**

Liver carboxylesterase 1, Acyl-coenzyme A:cholesterol acyltransferase, ACAT, Brain carboxylesterase hBr1, Carboxylesterase 1, CE-1, hCE-1, 3.1.1.1, Cocaine carboxylesterase, Egasyn, HMSE, Methylumbelliferyl-acetate deacetylase 1, 3.1.1.56, Monocyte/macrophage serine esterase, Retinyl ester hydrolase, REH, Serine esterase 1, Triacylglycerol hydrolase, TGH, CES1, CES2, SES1

Target/Specificity

Human CES1. This antibody is expected to recognise all three reported isoforms (NP_001020366.1; NP_001020365.1; NP_001257.4).

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

Carboxylesterase 1 / CES1 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

Carboxylesterase 1 / CES1 Antibody (Internal) - Protein Information**Name** CES1 ([HGNC:1863](#))**Synonyms** CES2, SES1**Function**

Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs (PubMed: [18762277](http://www.uniprot.org/citations/18762277)), (PubMed: [7980644](http://www.uniprot.org/citations/7980644)), (PubMed: [9169443](http://www.uniprot.org/citations/9169443)), (PubMed: [9490062](http://www.uniprot.org/citations/9490062)). Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty

acyl-CoA ester (PubMed:[18762277](http://www.uniprot.org/citations/18762277)), PubMed:[7980644](http://www.uniprot.org/citations/7980644), PubMed:[9169443](http://www.uniprot.org/citations/9169443), PubMed:[9490062](http://www.uniprot.org/citations/9490062)). Hydrolyzes the methyl ester group of cocaine to form benzoylcegonine (PubMed:[7980644](http://www.uniprot.org/citations/7980644)). Catalyzes the transesterification of cocaine to form cocaethylene (PubMed:[7980644](http://www.uniprot.org/citations/7980644)). Displays fatty acid ethyl ester synthase activity, catalyzing the ethyl esterification of oleic acid to ethyloleate (PubMed:[7980644](http://www.uniprot.org/citations/7980644)). Converts monoacylglycerides to free fatty acids and glycerol. Hydrolyzes of 2-arachidonoylglycerol and prostaglandins (PubMed:[21049984](http://www.uniprot.org/citations/21049984)). Hydrolyzes cellular cholesteryl esters to free cholesterol and promotes reverse cholesterol transport (RCT) by facilitating both the initial and final steps in the process (PubMed:[11015575](http://www.uniprot.org/citations/11015575), PubMed:[16024911](http://www.uniprot.org/citations/16024911), PubMed:[16971496](http://www.uniprot.org/citations/16971496), PubMed:[18762277](http://www.uniprot.org/citations/18762277)). First of all, allows free cholesterol efflux from macrophages to extracellular cholesterol acceptors and secondly, releases free cholesterol from lipoprotein-delivered cholesteryl esters in the liver for bile acid synthesis or direct secretion into the bile (PubMed:[16971496](http://www.uniprot.org/citations/16971496), PubMed:[18599737](http://www.uniprot.org/citations/18599737), PubMed:[18762277](http://www.uniprot.org/citations/18762277)).

Cellular Location

Endoplasmic reticulum lumen. Cytoplasm Lipid droplet. Note=Moves from cytoplasm to lipid droplets upon lipid loading. Associates with lipid droplets independently of triglycerides (TG) content of the droplets and hydrolyzes cholesteryl esters more efficiently from mixed droplets

Tissue Location

Expressed predominantly in liver with lower levels in heart and lung (PubMed:10562416).
Expressed in macrophages (PubMed:11015575, PubMed:18762277, PubMed:21049984)


Carboxylesterase 1 / CES1 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Carboxylesterase 1 / CES1 Antibody (Internal) - Images





Antibody (0.03 ug/ml) staining of Human Liver lysate (35 ug protein in RIPA buffer).



Anti-CES1 antibody IHC of human lung.

Carboxylesterase 1 / CES1 Antibody (Internal) - Background

Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl-CoA ester. Hydrolyzes the methyl ester group of cocaine to form benzoylecgonine. Catalyzes the transesterification of cocaine to form cocaethylene. Displays fatty acid ethyl ester synthase activity, catalyzing the ethyl esterification of oleic acid to ethyl oleate.

Carboxylesterase 1 / CES1 Antibody (Internal) - References

- Munger J.S., et al. J. Biol. Chem. 266:18832-18838(1991).
- Kroetz D.L., et al. Biochemistry 32:11606-11617(1993).
- Shibata F., et al. Genomics 17:76-82(1993).
- Becker A., et al. Arterioscler. Thromb. 14:1346-1355(1994).
- Islam M.R., et al. Arch. Biochem. Biophys. 372:53-61(1999).