

NR4A1 / NUR77 Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS12272**Specification**

NR4A1 / NUR77 Antibody (Internal) - Product Information

Application	IHC
Primary Accession	P22736
Reactivity	Human, Mouse, Rat, Rabbit, Hamster, Dog, Horse
Host	Goat
Clonality	Polyclonal
Calculated MW	64kDa KDa

NR4A1 / NUR77 Antibody (Internal) - Additional Information**Gene ID** 3164**Other Names**

Nuclear receptor subfamily 4 group A member 1, Early response protein NAK1, Nuclear hormone receptor NUR/77, Nur77, Orphan nuclear receptor HMR, Orphan nuclear receptor TR3, ST-59, Testicular receptor 3, NR4A1, GFRP1, HMR, NAK1

Target/Specificity

Human NR4A1 / NUR77. This antibody is expected to recognise isoform a (NP_002126.2 and NP_775180.1) only.

Format

TBS, pH 7.3, 0.02% Sodium Azide, 0.5% BSA

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

NR4A1 / NUR77 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

NR4A1 / NUR77 Antibody (Internal) - Protein Information**Name** NR4A1**Synonyms** GFRP1, HMR, NAK1**Function**

Orphan nuclear receptor. Binds the NGFI-B response element (NBRE) 5'-AAAGGTCA-3' (PubMed:18690216, PubMed:8121493, PubMed:9315652). Binds

9-cis-retinoic acid outside of its ligand- binding (NR LBD) domain (PubMed:18690216). Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation (PubMed:22983157). Regulates the inflammatory response in macrophages by regulating metabolic adaptations during inflammation, including repressing the transcription of genes involved in the citric acid cycle (TCA) (By similarity). Inhibits NF-kappa-B signaling by binding to low-affinity NF-kappa-B binding sites, such as at the IL2 promoter (PubMed:15466594). May act concomitantly with NR4A2 in regulating the expression of delayed-early genes during liver regeneration (By similarity). Plays a role in the vascular response to injury (By similarity).

Cellular Location

Nucleus. Cytoplasm, cytosol. Mitochondrion Note=Nuclear export to the cytosol is XPO1-mediated and positively regulated by IFI27 (PubMed:22427340). Translocation to the mitochondrion upon interaction with RXRA and upon the presence of 9-cis retinoic acid (PubMed:17761950).

Tissue Location

Fetal muscle and adult liver, brain and thyroid.

Volume


Array

NR4A1 / NUR77 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR4A1 / NUR77 Antibody (Internal) - Images



Anti-NUR77 antibody IHC of human breast.

NR4A1 / NUR77 Antibody (Internal) - Background

Orphan nuclear receptor. May act concomitantly with NURR1 in regulating the expression of delayed-early genes during liver regeneration. Binds the NGFI-B response element (NBRE) 5'-AAAAGGTCA-3' (By similarity). May inhibit NF-kappa-B transactivation of IL2. Participates in energy homeostasis by sequestering the kinase STK11 in the nucleus, thereby attenuating cytoplasmic AMPK activation.

NR4A1 / NUR77 Antibody (Internal) - References

Nakai A.,et al.Mol. Endocrinol. 4:1438-1443(1990).
Chang C.,et al.J. Steroid Biochem. 34:391-395(1989).
Ohkura N.,et al.Submitted (MAY-1996) to the EMBL/GenBank/DDBJ databases.
Kobayashi T.,et al.Submitted (DEC-2008) to the EMBL/GenBank/DDBJ databases.
Kaighin V.A.,et al.Submitted (DEC-2010) to the EMBL/GenBank/DDBJ databases.