

CD20 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS12340

Specification

CD20 Antibody (C-Terminus) - Product Information

Application	IHC
Primary Accession	P11836
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33kDa KDa

CD20 Antibody (C-Terminus) - Additional Information

Gene ID 931

Other Names

B-lymphocyte antigen CD20, B-lymphocyte surface antigen B1, Bp35, Leukocyte surface antigen Leu-16, Membrane-spanning 4-domains subfamily A member 1, CD20, MS4A1, CD20

Target/Specificity

Recognizes CD20. Species cross-reactivity: Human.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

CD20 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

CD20 Antibody (C-Terminus) - Protein Information

Name MS4A1

Synonyms CD20

Function

B-lymphocyte-specific membrane protein that plays a role in the regulation of cellular calcium influx necessary for the development, differentiation, and activation of B-lymphocytes (PubMed: [12920111](http://www.uniprot.org/citations/12920111), PubMed: [3925015](http://www.uniprot.org/citations/3925015), PubMed: [7684739](http://www.uniprot.org/citations/7684739)). Functions as a store-operated calcium (SOC) channel component promoting calcium influx after activation by the B-cell receptor/BCR (PubMed: [12920111](http://www.uniprot.org/citations/12920111), PubMed: [18474602](http://www.uniprot.org/citations/18474602), PubMed: [7684739](http://www.uniprot.org/citations/7684739)).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell membrane; Lipid-anchor. Note=Constitutively associated with membrane rafts.

Tissue Location


Expressed on B-cells.

CD20 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD20 Antibody (C-Terminus) - Images



Anti-MS4A1 / CD20 antibody IHC of human spleen.

CD20 Antibody (C-Terminus) - Background

This protein may be involved in the regulation of B-cell activation and proliferation.

CD20 Antibody (C-Terminus) - References

- Stamenkovic I., et al. *J. Exp. Med.* 167:1975-1980(1988).
Tedder T.F., et al. *Proc. Natl. Acad. Sci. U.S.A.* 85:208-212(1988).
Einfeld D.A., et al. *EMBO J.* 7:711-717(1988).
Tedder T.F., et al. *J. Immunol.* 142:2560-2568(1989).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).