

**S100A4 / FSP1 Antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS12477****Specification**

---

**S100A4 / FSP1 Antibody - Product Information**

Application	IHC
Primary Accession	<a href="#">P26447</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	12kDa KDa

**S100A4 / FSP1 Antibody - Additional Information****Gene ID** 6275**Other Names**

Protein S100-A4, Calvasculin, Metastasin, Placental calcium-binding protein, Protein Mts1, S100 calcium-binding protein A4, S100A4, CAPL, MTS1

**Reconstitution & Storage**

+4°C or -20°C, Avoid repeated freezing and thawing.

**Precautions**

S100A4 / FSP1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**S100A4 / FSP1 Antibody - Protein Information****Name** S100A4**Synonyms** CAPL, MTS1**Function**

Calcium-binding protein that plays a role in various cellular processes including motility, angiogenesis, cell differentiation, apoptosis, and autophagy (PubMed: [16707441](http://www.uniprot.org/citations/16707441), PubMed: [23752197](http://www.uniprot.org/citations/23752197), PubMed: [30713770](http://www.uniprot.org/citations/30713770)). Increases cell motility and invasiveness by interacting with non-muscle myosin heavy chain (NMMHC) IIA/MYH9 (PubMed: [16707441](http://www.uniprot.org/citations/16707441)). Mechanistically, promotes filament depolymerization and increases the amount of soluble myosin-IIA, resulting in the formation of stable protrusions facilitating chemotaxis (By similarity). Modulates also the pro-apoptotic function of TP53 by binding to its C-terminal transactivation domain within the nucleus and reducing its protein levels (PubMed: [23752197](http://www.uniprot.org/citations/23752197)). Within the extracellular space, stimulates cytokine production including granulocyte colony-

stimulating factor and CCL24 from T-lymphocytes (By similarity). In addition, stimulates T-lymphocyte chemotaxis by acting as a chemoattractant complex with PGLYRP1 that promotes lymphocyte migration via CCR5 and CXCR3 receptors (PubMed:<a href="http://www.uniprot.org/citations/26654597" target="\_blank">26654597</a>, PubMed:<a href="http://www.uniprot.org/citations/30713770" target="\_blank">30713770</a>).

#### Cellular Location

Secreted. Nucleus Cytoplasm {ECO:0000250|UniProtKB:P07091}

#### Tissue Location


Ubiquitously expressed.

### S100A4 / FSP1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### S100A4 / FSP1 Antibody - Images



Anti-S100A4 antibody IHC of human small intestine.

### S100A4 / FSP1 Antibody - References

Engelkamp D.,et al.Biochemistry 31:10258-10264(1992).  
Engelkamp D.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:6547-6551(1993).  
Tulchinsky E.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:9146-9150(1992).  
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).